

UNIVERSIDADE FEDERAL DE ALFENAS

KAREN CRISTINA OLIVEIRA

**EPITOPE-BASED VACCINE OF A *Brucella abortus* PUTATIVE SMALL RNA
TARGET INDUCES PROTECTION AND LESS TISSUE DAMAGE IN MICE**

ALFENAS/MG

2021

KAREN CRISTINA OLIVEIRA

**EPITOPE-BASED VACCINE OF A *Brucella abortus* PUTATIVE SMALL RNA
TARGET INDUCES PROTECTION AND LESS TISSUE DAMAGE IN MICE**

Dissertação apresentada como parte dos requisitos exigidos para obtenção do título de Mestre pelo Programa de Pós-Graduação em Ciências Biológicas da Universidade Federal de Alfenas - MG. Área de Concentração: Biologia Celular, Molecular e Estrutural das Doenças Agudas e Crônicas.

Orientador: Prof. Dr. Leonardo Augusto de Almeida

ALFENAS/MG

2021

Dados Internacionais de Catalogação-na-Publicação (CIP)
Sistema de Bibliotecas da Universidade Federal de Alfenas
Biblioteca Central – Campus Sede

Oliveira, Karen Cristina
O48e Epitope-based vaccine of a *Brucella abortus* putative small rna target induces protection and less tissue damage in mice. / Karen Cristina Oliveira – Alfenas, MG, 2021.
[51] f.: il. –

Orientador: Leonardo Augusto de Almeida.
Dissertação (Mestrado em Ciências Biológicas) – Universidade Federal de Alfenas, 2021.
Bibliografia.

1. *Brucella abortus*. 2. Vacina. 3. Resposta imune. 4. Vacinologia reversa. 5. Brucelose. 6. Apolipoproteína N-aciltransferase. I. Almeida, Leonardo Augusto de. II. Título.

CDD- 570

KAREN CRISTINA OLIVEIRA

EPITOPE-BASED VACCINE OF A *Brucella abortus* PUTATIVE SMALL RNA TARGET INDUCES PROTECTION AND LESS TISSUE DAMAGE IN MICE

A Banca examinadora abaixo-assinada aprova a Dissertação apresentada como parte dos requisitos para a obtenção do título de Mestra em Ciências Biológicas pela Universidade Federal de Alfenas. Área de concentração: Biologia Celular, Molecular e Estrutural das Doenças Agudas e Crônicas.

Aprovada em: 23 de setembro de 2021.

Prof. Dr. Leonardo Augusto de Almeida
Instituição: Universidade Federal de Alfenas - UNIFAL-MG

Prof. Dr. Gilson Costa Macedo
Instituição: Universidade Federal de Juiz de Fora - UFJF

Prof. Dr. Fábio Antônio Colombo
Instituição: Universidade Federal de Alfenas - UNIFAL-MG



Documento assinado eletronicamente por **Leonardo Augusto de Almeida, Professor do Magistério Superior**, em 24/09/2021, às 10:22, conforme horário oficial de Brasília, com fundamento no art. 6º, § 1º, do [Decreto nº 8.539, de 8 de outubro de 2015](#).



Documento assinado eletronicamente por **Fabio Antonio Colombo, Professor do Magistério Superior**, em 24/09/2021, às 13:28, conforme horário oficial de Brasília, com fundamento no art. 6º, § 1º, do [Decreto nº 8.539, de 8 de outubro de 2015](#).



Documento assinado eletronicamente por **Gilson Costa Macedo, Usuário Externo**, em 27/09/2021, às 12:04, conforme horário oficial de Brasília, com fundamento no art. 6º, § 1º, do [Decreto nº 8.539, de 8 de outubro de 2015](#).



A autenticidade deste documento pode ser conferida no site https://sei.unifal-mg.edu.br/sei/controlador_externo.php?acao=documento_conferir&id_orgao_acesso_externo=0, informando o código verificador **0597579** e o código CRC **4951506B**.

Dedico este trabalho à minha mãe,
meus irmãos Karla e Cadú e à vovó
Nilza (*in memoriam*).

AGRADECIMENTOS

À Deus, pela saúde, força, persistência e resiliência concedidas a mim durante esse período para que fosse possível a conclusão mais uma etapa!

Ao professor Leonardo Augusto de Almeida pela orientação e profissionalismo. Agradeço pela paciência e confiança, contribuindo para com o meu amadurecimento, enriquecimento e crescimento profissional, despertando ainda mais o meu interesse pela pesquisa. Obrigada por toda a dedicação, pelos ensinamentos e por me acolher como sua orientada.

Aos amigos do Labiomol, em especial á Natália, Thiago, Ana, Bianca, Jéssica, Caio e Gustavo, por todos os momentos que compartilhamos (inclusive a parceria nas experimentações e as fugidas pra ir na Don'ana rrsr), fazendo com que os dias se tornassem mais alegres, leves e divertidos. Obrigada por todo o apoio, colaboração, amizade e risadas. Levarei vocês no coração pro resto da vida!

À professora Patrícia Paiva Corsetti e seus orientados Evandro, Thaís e Ana Maria, pelas trocas de conhecimentos e por terem se tornado meus amigos.

Ao Programa de Pós-Graduação em Ciências Biológicas, professores - por todo o conhecimento transmitido, Martha - pela convivência nas disciplinas e durante as reuniões do colegiado e todos os técnicos e funcionários do Departamento de Microbiologia e Imunologia da Unifal: obrigada a todos vocês.

Aos meus familiares, em especial minha Mãe Joelma, meu pai Carlos e meus irmãos Karla e Cadu pelo amor, carinho, incentivo e principalmente por acreditarem no meu sonho.

Ao meu namorado Thiago pela paciência, incentivo, carinho e amor. Obrigada por aturar minhas reclamações, estresses, e principalmente por ser a pessoa que mais acreditou em mim e nos meus sonhos (sonhando e caminhando sempre ao meu lado).

Aos meus quatro melhores amigos da vida, Weverton, Daniel, Fernanda e Taís por estarem sempre do meu lado, fortalecendo minha base e acreditando nos meus sonhos.

À Universidade Federal de Alfenas e ao Programa de Pós-Graduação em Ciências Biológicas pela oportunidade e pelo meu crescimento acadêmico e à todos que contribuíram direta ou indiretamente com o desenvolvimento deste trabalho.

Agradeço, de antemão, a disponibilidade de todos que compuseram minha banca de defesa da dissertação: Prof. Dr. Gilson Costa Macedo, Prof. Dr. Fábio Antônio

Colombo, Prof. Dr. Eduardo de Figueiredo Peloso, Prof. Dra. Valéria Quintana Cavicchioli e, novamente, ao Prof. Dr. Leonardo Augusto de Almeida.

À CAPES pela concessão da bolsa de estudos.

O presente trabalho foi realizado com apoio da Coordenação de Aperfeiçoamento de Pessoal de Nível Superior - Brasil (CAPES) – Código de Financiamento 001.

“Vocês jovens, doutores, cientistas do futuro, não se deixem abalar por um ceticismo estéril, nem se deixem desencorajar pela tristeza de certas horas que as nações passam.”

(Louis Pasteur, 1865)

RESUMO

Brucella abortus é uma bactéria intracelular Gram-negativa que causa uma doença zoonótica chamada brucelose. Embora as vacinas atualmente disponíveis para a imunização animal possam ter potencial imunogênico, essas ainda apresentam muitas desvantagens, causando abortos de grande proporção em fêmeas prenhas e febre ondulante em humanos. Neste contexto, a recente tendência no projeto de novas vacinas contra brucelose, têm se baseado na estratégia de predição de epítomos imunogênicos selecionados por vacinologia reversa. Sendo assim, o objetivo deste estudo foi identificar e avaliar a imunogenicidade de um epítomo vacinal alvo de putativos pequenos RNAs de *B. abortus* mediante a infecção desta bactéria em modelo murino. Foi demonstrado nesse trabalho que pequenos RNAs de *B. abortus* são expressos durante a infecção precoce de macrófagos derivados da medula óssea (BMDMs), sendo identificado uma apolipoproteína N-aciltransferase (Int) como o putativo alvo de maior expressão dos pequenos RNAs. Visto que a apolipoproteína N-aciltransferase apresenta diminuição da sua expressão em modelo de BMDMs infectados, um epítomo desta proteína foi racionalmente selecionado por imunoinformática e explorado como candidato a vacinação contra brucelose. Camundongos C57BL/6 imunizados e desafiados com *B. abortus* mostraram menor recuperação no número de bactérias viáveis no fígado, baço e linfonodo axilar quando comparados a camundongos não vacinados. Os camundongos vacinados e infectados apresentaram aumento na expressão de TNF- α , IFN- γ e IL-6, seguido do também aumento na expressão dos genes anti-inflamatórios IL-10 e TGF- β no fígado, justificando a redução no número e tamanho dos granulomas observados. BMDMs estimulados com sobrenadante de esplenócitos de camundongos vacinados e infectados apresentaram marcação para CD86+ mais intensa que os demais estímulos, além de expressarem maior quantidade de iNOS e consequente aumento na produção de NO, sugerindo aumento na capacidade fagocítica e microbicida dessas células em eliminar a bactéria. Em conjunto, os resultados demonstraram que o peptídeo vacinal foi capaz de estimular uma resposta imune protetora em organismos infectados com características sugestivas de predominância do perfil Th1.

Palavras-chave: *Brucella abortus*; Vacina; Resposta imune; Vacinologia reversa; Brucelose; Apolipoproteína N-aciltransferase.

ABSTRACT

Brucella abortus is a Gram-negative intracellular bacterium that causes a zoonotic disease called brucellosis. Although currently available vaccines for animal immunization have immunogenic potential, they still have many disadvantages, causing large-scale abortions in pregnant females and undulating fever in humans. In this context, the recent trend in the design of new vaccines against brucellosis has been based on the strategy of prediction of immunogenic epitopes selected by reverse vaccinology. Therefore, this study aimed to identify and evaluate the immunogenicity of a target vaccine epitope of putative small RNAs of *B. abortus* upon infection of this bacterium in a murine model. It was demonstrated in this work that small RNAs of *B. abortus* are expressed during the early infection of bone marrow-derived macrophages (BMDMs), and an apolipoprotein N-acyltransferase (Int) was identified as the putative target of higher expression of small RNAs. Since the apolipoprotein N-acyltransferase has decreased expression in a model of infected BMDMs, an epitope of this protein was rationally selected by immunoinformatics and explored as a candidate for vaccination against brucellosis. C57BL/6 mice immunized and challenged with *B. abortus* showed lower recovery in the number of viable bacteria in the liver, spleen, and axillary lymph node when compared to non-vaccinated mice. The vaccinated and infected mice showed an increase in the expression of TNF- α , IFN- γ , and IL-6, followed by an increase in the expression of the anti-inflammatory genes IL-10 and TGF- β in the liver, justifying the reduction in number and size of the observed granulomas. BMDMs stimulated with supernatant from splenocytes from vaccinated and infected mice showed more intense CD86⁺ marking than the other stimuli, in addition to expressing a greater amount of iNOS and consequent increase in NO production, suggesting an increase in the phagocytic and microbicidal capacity of these cells to eliminate the bacteria. Together, the results demonstrated that the vaccine peptide was able to stimulate a protective immune response in infected organisms with characteristics suggestive of a predominance of the Th1 profile.

Keywords: *Brucella abortus*; vaccine; immune response; reverse vaccinology; brucellosis; apolipoprotein N-acyltransferase.

LISTA DE ABREVIATURAS E SIGLAS

BMDM	Bone Marrow-Derived Macrophages
CEUA	Committee on the Ethics of Animal Experiments
CFU	Colony Forming Units
FBS	Fetal Bovine Serum
HLA	Human Leukocyte Antigen
IEDB	Immune Epitope Database
IL	Interleukin
LPS	Lipopolysaccharide
MHC	Major histocompatibility complex
NCBI	National Center for Biotechnology Information
NO	Nitric Oxid
PBS	Phosphate-buffered saline
RV	Reverse vaccinology
SLC	Subcellular localization
TGF	Transforming Growth Factor
TNF	Tumor necrosis factor

SUMÁRIO

1 INTRODUÇÃO.....	11
CAPÍTULO 1	14
ARTIGO: Epitope-based vaccine of a <i>Brucella abortus</i> putative small RNA target induces protection and less tissue damage in mice	14
2 CONCLUSÃO.....	15
REFERÊNCIAS	16

1 INTRODUÇÃO

O gênero *Brucella* compreende bactérias que apresentam o formato de cocobacilos Gram-negativos, imóveis, não esporogênicos e intracelulares facultativos (TANA *et al.*, 2021). Estas bactérias são responsáveis por causar a brucelose, uma zoonose negligenciada amplamente distribuída no mundo, em que qualquer órgão ou tecido do organismo pode ser acometido com incidência variável (JAMIL *et al.*, 2020; TIAN *et al.*, 2020). Apesar dos esforços feitos para controlar a doença em muitos países (ELFAKI *et al.*, 2015), as transmissões dadas por *Brucella* persistem em animais domésticos e, conseqüentemente, infecções ocorrem frequentemente em humanos. As bactérias do gênero *Brucella* exibem características antigênicas e metabólicas distintas, e são classificadas de acordo com sua preferência pelo hospedeiro animal. Este gênero, atualmente, abrange doze espécies isoladas e nomeadas (HULL, SCHUMAKER, 2018), dentre as quais *B. melitensis*, *B. abortus* e *B. suis* são os principais agentes etiológicos da brucelose humana (EL-SAYED, AWAD, 2018), sendo a *B. melitensis* a mais patogênica e a *B. abortus* a mais difundida no mundo, uma vez que a doença atinge anualmente cerca de 500 mil pessoas em todo o mundo (GŁOWACKA *et al.*, 2018). *B. abortus* têm como hospedeiro preferencial os bovinos e mensura-se que 300 milhões das 1,4 bilhões cabeças de gado do mundo estejam infectadas com *B. abortus* (ELFAKI *et al.*, 2015; O'CALLAGHAN, 2020).

As principais formas de transmissão de *B. abortus* para os homens é a partir da ingestão de lácteos contaminados não submetidos ao processo de pasteurização (LINDAHL-RAJALA *et al.*, 2017), carne crua proveniente de animais infectados (CASALINUOVO *et al.*, 2016), pela inalação de aerossóis que contêm o patógeno. Já, entre os animais a contaminação se dá por contato direto com tecidos animais infectados e suas secreções, e pelo hábito dos bovinos de lamber e cheirar animais recém-nascidos ou mesmo fetos abortados (POESTER *et al.*, 2013). A brucelose é uma doença sistêmica na qual qualquer órgão ou tecido do organismo pode estar envolvido. Nos animais a brucelose é uma infecção crônica, que pode persistir por toda a vida do hospedeiro. Nas fêmeas, *Brucella* possui tropismo pelo hormônio placentário bovino, o eritritol, levando a lesões nas glândulas uterinas enquanto que, nos machos, a bactéria possui tropismo por hormônios masculinos, como a testosterona, dirigindo-se aos testículos (DORNELES *et al.*, 2015; GUTIÉRREZ-JIMÉNEZ *et al.*, 2018). Assim, a infecção por *B. abortus* afeta principalmente os órgãos reprodutivos causando aborto e infertilidade. A brucelose

humana raramente é fatal, porém é uma doença severamente debilitante e incapacitante que apresenta tendência à cronicidade e persistência (BYNDLOSS, TSOLIS, 2016; HASANJANI, EBRAHIMPOUR, 2015).

Por ser um microrganismo intracelular com tropismo principalmente por células do sistema mononuclear fagocitário, a resposta à *Brucella* spp. envolve toda a gama do sistema imunológico, desde a imunidade inata à imunidade adaptativa. A resposta protetora eficaz contra a infecção por *B. abortus* requer a ativação de linfócitos TCD4+ e TCD8+, produção de citocinas e produtos do perfil de resposta Th1 característico contra bactérias intracelulares, como IFN- γ e TNF- α , além da ativação de macrófagos e células dendríticas (JEZI *et al.*, 2019). Considerando os mecanismos de virulência de *B. abortus*, aliados ao fato da bactéria estar localizada intracelularmente, a antibioticoterapia torna-se particularmente pouco eficaz nesses casos (GLOWACKA *et al.*, 2018). Levando em consideração tais aspectos, a vacinação dos animais caracteriza a principal medida preventiva com capacidade de conferir proteção efetiva e duradoura contra a infecção por *B. abortus* (CARVALHO *et al.*, 2020).

No mercado veterinário atual, as cepas S19 (linhagem lisa) e RB51 (linhagem rugosa) têm sido usadas para controlar brucelose bovina em todo o mundo. No entanto, ambas cepas apresentam riscos potenciais para os animais e humanos (DE OLIVEIRA *et al.*, 2021). Nos animais, a vacinação utilizando as cepas atenuadas vivas S19 e RB51 pode ocasionar a reversão de virulência, provocando abortos, artropatias, orquite e infertilidade, além de induzir a produção de anticorpos, o que interfere no diagnóstico sorológico da doença (GRUPTA *et al.*, 2020; LALSIAMTHARA, LEE, 2017). Ademais, ambas as vacinas são secretadas no leite e podem infectar humanos em contato direto e causar abortos em animais grávidas (FRANC *et al.*, 2018). Para humanos, a exposição acidental às vacinas contra brucelose caracteriza uma importante fonte de contaminação, uma vez que essas são patogênicas, podendo causar diversas morbidades, dentre elas: febre ondulante provocada devido a episódios de bacteremia seguido por novo foco de infecção, artrite, endocardite, osteomielite e complicações neurológicas (BALDI, GIAMBARTOLOMEI, 2013; FRANC *et al.*, 2018). Até então, não há nenhuma vacina destinada a imunização humana contra brucelose, apenas animal.

Atualmente, objetivando desenvolver uma vacina segura e melhorada contra a brucelose, numerosos estudos tem sido realizados para entender os mecanismos de imunidade protetora de *Brucella* no modelo murino (CARVALHO *et al.*, 2016; GHASEMI *et al.*, 2014) e também nos hospedeiros naturais (NOL *et al.*, 2016; OLSEN

et al., 2015). Nesse contexto, a vacinologia reversa tem sido utilizada com uma importante ferramenta moderna capaz de predizer *in silico* antígenos de *Brucella* com propriedades imunogênicas, capazes de promover resposta humoral e celular com produção de citocinas de resposta Th1, Th2 e Th17 (GOMEZ *et al.*, 2013), e permitindo o estudo de novas vacinas de *Brucella* (VISHNU *et al.*, 2015). A aquisição de conhecimentos genômicos, proteômicos, tecnologias em vacina e técnicas de DNA recombinante tem possibilitado um número crescente de pesquisas com vacinas mais seguras (ESCALONA *et al.*, 2017), entretanto, apenas alguns antígenos isolados demonstram imunidade protetora importante *in vivo* (ABKAR *et al.*, 2015; JAIN *et al.*, 2014). Sabe-se que as respostas imunológicas do tipo Th1 contra a infecção de *Brucella* são melhores estimuladas por vacinas de cepas vivas atenuadas ou cepas mutantes (CARVALHO *et al.*, 2016). Embora a vacinologia reversa seja um método ainda não aplicado em vacinas comerciais contra *Brucella* spp., a mesma provou ser útil na identificação de antígenos protetores contra outros patógenos importantes, por exemplo, o meningococo do sorogrupo B (PIZZA *et al.*, 2000), caracterizando a eficácia dessa ferramenta.

Dada a relevância da bactéria *B. abortus* sendo causadora de uma zoonose de importância médico-sanitária e estimulados pela necessidade de novas estratégias vacinais mais seguras, o objetivo deste trabalho foi identificar e avaliar a imunogenicidade de um epítipo vacinal alvo de putativos pequenos RNAs de *B. abortus* mediante a infecção desta bactéria em modelo murino.

CAPÍTULO 1

ARTIGO: Epitope-based vaccine of a *Brucella abortus* putative small RNA target induces protection and less tissue damage in mice

1 **Epitope-based vaccine of a *Brucella abortus* putative small RNA target induces**
2 **protection and less tissue damage in mice**

3 **Karen Cristina Oliveira¹, Gustavo Andrade Brancaglioni¹, Natália Cristina de Melo Santos¹,**
4 **Leonardo Pereira de Araújo¹, Evandro Novaes², Renato Lima Santos³, Sergio Costa**
5 **Oliveira⁴, Patrícia Paiva Corsetti^{1*}, Leonardo Augusto de Almeida^{1*}**

6 ¹Laboratory of Molecular Biology of Microorganisms, Federal University of Alfenas, Alfenas,
7 Minas Gerais, Brazil.

8 ²Department of Biology, Federal University of Lavras, Lavras, Minas Gerais, Brazil.

9 ³Veterinary School, Department of Clinic and Veterinary Surgery, Federal University of Minas
10 Gerais, Belo Horizonte, Minas Gerais, Brazil.

11 ⁴Department of Biochemistry and Immunology, Federal University of Minas Gerais, Belo
12 Horizonte, Minas Gerais, Brazil.

13 ***Correspondence:**

14 Leonardo Augusto de Almeida

15 E-mail: leonardo.almeida@unifal-mg.edu.br

16 Patrícia Paiva Corsetti

17 E-mail: ppcorsetti@hotmail.com

18

19 **Abstract**

20 *Brucella* spp. are Gram-negative, facultative intracellular bacteria that cause brucellosis in humans
21 and animals. Currently available live attenuated vaccines against brucellosis still have drawbacks.
22 Therefore, subunit vaccines, produced using epitope-based antigens, have the advantage of being
23 safe, cost-effective and efficacious. Here, we identified *B. abortus* small RNAs expressed during
24 early infection with bone marrow-derived macrophages (BMDMs) and an apolipoprotein N-
25 acyltransferase (Int) was identified as the putative target of the greatest expressed small RNA.
26 Decreased expression of Int was observed during BMDM infection and the protein sequence was
27 evaluated to rationally select a putative immunogenic epitope by immunoinformatic, which was
28 explored as a vaccinal candidate. C57BL/6 mice were immunized and challenged with *B. abortus*,
29 showing lower recovery in the number of viable bacteria in the liver, spleen, and axillary lymph
30 node when compared to non-vaccinated mice. The vaccinated and infected mice showed the
31 increased expression of *TNF- α* , *IFN- γ* , and IL-6 following expression of the anti-inflammatory
32 genes *IL-10* and *TGF- β* in the liver, justifying the reduction in the number and size of the observed
33 granulomas. BMDMs stimulated with splenocyte supernatants from vaccinated and infected mice
34 increase the CD86+ marker, as well as expressing greater amounts of iNOS and the consequent
35 increase in NO production, suggesting an increase in the phagocytic and microbicidal capacity of
36 these cells to eliminate the bacteria.

37 **Keywords:** *Brucella abortus*; vaccine; immune response; reverse vaccinology; brucellosis;
38 apolipoprotein N-acyltransferase

39 **1 Introduction**

40 Brucellosis is a global zoonotic infectious disease caused by bacteria of the genus *Brucella*.
41 The disease is a serious public health threat worldwide, particularly in developing countries of
42 Central Asia, Africa, South America, and the Mediterranean region (1). Brucellosis affects
43 mammals, causing abortion and infertility in affected animals. This infection can spread from
44 animals to humans, mainly via the ingestion of unpasteurized milk or dairy products and, to a lesser
45 extent, via direct contact with infected animals (2). In humans, brucellosis can cause a severe febrile
46 disease with various clinical complications ranging from mild to severe symptoms including
47 undulant fever, joint pain arthritis, endocarditis, and meningitis (3-5). The genus *Brucella* includes
48 Gram-negative facultative intracellular bacteria from Alphaproteobacteria, and, currently, the
49 genus consists of 12 species that are classified based on their host preferences (6). Although
50 several *Brucella* species are potentially zoonotic agents, *Brucella melitensis*, *Brucella abortus*,
51 and *Brucella suis* are considered the most pathogenic *Brucella* species that have a serious impact
52 on public health and the livestock industry (7,8), with *B. abortus* being the most widespread
53 throughout the world, according to the World Health Organization (WHO) (9,10). Since brucellosis
54 is the most common zoonotic disease worldwide and has become a serious concern in recent years
55 (11), the strategy used to control brucellosis depends mainly on the massive vaccination of domestic
56 animals to prevent the disease from spreading to healthy animals and humans (12,13). Almost all
57 vaccines against *Brucella* spp. are live attenuated strains with extensive global use but with various
58 drawbacks, such as pathogenicity to humans and residual virulence in animals, which can cause
59 abortion, orchitis, and infertility (14-16). Moreover, it is difficult to differentiate infected animals
60 from vaccinated animals by serological tests. These drawbacks have prompted several research
61 groups to attempt the development of safer vaccines.

62 Subunit vaccines have promising applications with the advantage of being safe, cost-
63 effective, and efficacious. During the past two decades, various antigens have been extracted
64 from *Brucella*, such as Omp19, Omp25, L7/L12, P39, SodC, InpB, AsnC, and TF (17-24). These
65 available antigens have been shown to provide protection against *Brucella* infection by reducing
66 the organ's bacterial load in mice. While such findings are highly promising, subunit vaccines
67 using known antigens cannot provide the levels of protection conferred by live attenuated vaccines
68 (25). Further investigation is needed to identify novel antigens, and increase vaccine efficacy. In
69 contrast to the conventional vaccine development that requires cultivation and extensive empirical
70 screening, reverse vaccinology (RV) is an interesting *in silico* approach to identify protective
71 antigens using pathogen genomic data (26-29). RV has been implemented to identify protective
72 antigens of numerous pathogens, including *B. abortus* (30-32).

73 The main goal of this study was to screen potential antigens in the genome of *B. abortus*
74 using RV as a search strategy and subsequently evaluating the immunogenic capacity of the
75 peptide in an animal model. We used an *in silico* methodology to select epitopes candidates based
76 on their biological characteristics strongly associated with protective antigenicity from putative
77 targets of small RNAs expressed in infected BMDMs. From these predictions, a transmembrane
78 epitope of apolipoprotein N-acyltransferase was selected for efficacy verification in a mouse
79 model showing promising results to be used as an epitope-based vaccine against brucellosis that
80 may induce robust immunity against the bacterium.

81 **2 Material and methods**

82 **2.1 Ethics statement**

83 This study was carried in strict accordance with the Brazilian laws 6638 and 9605 in Animal
84 Experimentation. The protocol was approved by the Committee on the Ethics of Animal
85 Experiments of the Federal University of Alfenas (CEUA 16/2020).

87 **2.2 Mice, cell culture and bacteria**

88 The strain C57BL/6 mice aged 6–8 weeks were purchased from the Federal University of Minas
89 Gerais animal facility (UFMG, Belo Horizonte, Brazil). Bone marrow cells were obtained from
90 femora and tibia of mice and they were grown in bone marrow-derived macrophages (BMDMs) as

91 previously described by our group [33]. *B. abortus* virulent strain 2308 was obtained from our own
92 laboratory collection. They were grown in *Brucella* broth medium (BD Pharmingen, San Diego,
93 CA, USA) for 3 days at 37°C.

94 95 **2.3 BMDM infection with *B. abortus***

96 BMDMs were infected with virulent *B. abortus* strain 2308 at a multiplicity of infection of 100:1.
97 Bacteria were centrifuged onto macrophages at 400 × g for 10 min at 4°C and then cells were
98 incubated for 30 min at 37°C under 5% CO₂. Macrophages were extensively washed with HBSS to
99 remove extracellular bacteria and incubated for an additional 90 min in medium supplemented with
100 100 µg/mL gentamycin to kill extracellular bacteria. Thereafter, the antibiotic concentration was
101 decreased to 10 µg/mL. Thirty minutes after infection, BMDMs were washed three times with
102 HBSS before processing following homogenization with 100µl of LS TRIzol® reagent Invitrogen
103 (Waltham, Massachusetts, EUA) for total RNA isolation.

104 105 **2.4 Small RNA sequencing and bioinformatics identification**

106 The construction and sequencing of a strand-specific small RNA (15–50 nt) library was conducted
107 by FASTERIS SA (Plan-les-Ouates, Switzerland), based on the Illumina® TruSeq® Small RNA
108 Library Prep Kit for Illumina HiSeq 2000 sequencing (Illumina Inc., San Diego, CA, USA). To
109 remove adapter sequences from small RNA raw reads of sequencing, the Cutadapt tool was used
110 [34] and sequencing quality was analyzed using the Trimmomatic V0.32 tool (35). Reads were
111 mapped to the *B. melitensis* biovar *Abortus* (strain 2308) genome using the Bowtie program (36) to
112 report the best alignment for each read allowing a maximum of one replacement per alignment.
113 Using BedTool, reads mapped at alignment were analyzed to determine the depth of coverage
114 across the genome determining the hotspots areas of sRNAs (37). Coverage with more than 50
115 reads and areas with a distance of less than 50bp were joined as the same hotspot area.

116 117 **2.5 Bioinformatic and reverse vaccinology**

118 To assess the putative targets of those small RNAs highly expressed by *B. abortus* during infection,
119 one hundred coding genes were obtained using the intersect BedTool to indicate the overlap
120 between small RNAs and coding genes (37). As the overlaps showed the possibility of affecting
121 more than 50% of the *B. abortus* coding genes, which genes contain the hottest hotspots (i.e., with
122 the highest average coverage) was evaluated. The proteins from the putative target coding genes
123 were filtered by assigning selection criteria according to biological characteristics: (1) subcellular
124 location (SCL) using Psortb v3.0.2 program (38), (2) the presence of a signal peptide using SignalP
125 5.0 Server tool (39), and (3) the presence of transmembrane helices and exposed regions using three
126 tools: TMPRED, SOSUI 1.11, and TMHMM 2.0 (40-42). The obtained proteins were then analyzed
127 for biological function using an online tool UniProt Knowledgebase (UniProtKB) and only one of
128 the 7 was selected [43]. The selected protein apolipoprotein N-acyltransferase (WP_002965220.1)
129 was used to identify epitopes composed of fifteen amino acids present in the extracellular portion
130 of Int. The extracellular portion of the surface-associated target protein was subjected to the
131 sequential mapping of epitopes predicted to bind tightly to major histocompatibility complex
132 (MCH) class II, using the tools NetMHCII 2.2, SYFPEITHI, and RANKPEP (44-46). To obtain the
133 best candidates of putative immunogenic epitopes from Int, using Multalin 5.4, the obtained
134 epitopes were aligned, and the linear amino acid sequence including approximately 15 amino acids
135 that presented a repeats sequence was selected (47). For the epitope antigenicity test, the VaxiJen
136 2.0 server was used with the 0.5 and “probable antigen” cutoff (48). For the allergenicity test, the
137 server “AlgPred: Prediction of Allergenic Proteins and Mapping of IgE Epitopes” was used with
138 the hybrid method that consists of using the following five tools available on the server: SVMc,
139 IgE, epitope, ARPs BLAST, and MAST (49). For the epitope similarity analysis, the Protein
140 BLAST tool available on the NCBI platform was used (50). For the physical and chemical
141 properties test, the ExPASy server ProtParam tool was used with a cutoff of 40 for structure stability
142 values (51). The pipeline of bioinformatic analysis is depicted and summarized in Figure 1.

2.6 Real-time RT-PCR for apolipoprotein N-acyltransferase expression

144 Thirty-minute *B. abortus*-infected BMDMs or the exponential growth of *B. abortus* in *Brucella*
145 broth medium total RNA were extracted using TRIzol reagent (Invitrogen). Reverse-transcription
146 of 100 ng from total RNA was performed using random primers according to the Illustra™ Ready-
147 To-Go RT-PCR Beads kit (GE Healthcare, Buckinghamshire, UK). Real-time RT-PCR was
148 conducted in a final volume of 10 µL containing the following: SYBR® Green PCR Master Mix
149 (Applied Biosystems, Foster City, CA, USA), with cDNA as the PCR template and primers to
150 amplify specific fragments corresponding to specific gene targets: IntF: 5`-
151 CTGATGTGATTGTCTGGCCG-3`, IntR: 5`-CCTGAGGTGTCGATTCCAGT-3`, *Brucella* 16S
152 F: 5`- TCTCACGACACGAGCTGACG -3`, *Brucella* 16S R: 5`- CCTGAGGTGTCGATTCCAGT
153 -3`. The PCR reaction was performed using the ABI 7500 Real-Time PCR System (Applied
154 Biosystems, Foster City, CA), with the following cycling parameters: 60°C for 10 min, 95°C for
155 10 min, 40 cycles of 95°C for 15 sec and 60°C for 1 min, and a dissociation stage of 95°C for 15
156 sec, 60°C for 1 min, 95°C for 15 sec, and 60°C for 15 sec. All data are presented as relative
157 expression units after normalization to the *Brucella* 16S gene. PCR measurements were conducted
158 in triplicate. The differences in the relative expression were analyzed by Student's *t* test with a two-
159 tailed distribution ($p < 0.05$ indicates statistical significance).
160
161

2.7 Int structural modeling and epitope-MHCII docking

162 The Int protein from *Brucella* was retrieved via the NCBI database, followed by molecular
163 modeling by homology via WebServer Phyre², which uses the Markov model for the best possible
164 global alignments to generate the most accurate protein in its main function (52). After modeling,
165 the epitope under investigation from the protein was obtained, and the preparation was performed
166 using the MGL tool, in which missing atoms were corrected and water molecules were removed
167 (53). The same preparation was performed for MHCII molecules found on the PDB server (Protein
168 Data Bank) (54). For docking execution, both files were converted to the PDBQT format required
169 by AutoDock Vina (55), using the Openbabel tool (56). The gridbox was generated around the
170 active sites of the recovered MHCII proteins. To visualize the interactions between epitope-MHCII
171 in 2D and 3D diagram format, the Ligplot+, and Pymol tools were used, respectively (57,58).
172
173

2.8 Mice experiments

174 To assess the ability of vaccinal peptide to induce immune responses, 6–8-week-old C57BL/6 mice
175 were contained in cages on a 12:12 light/dark cycle and fed *ad libitum* with standard rodent diet
176 and no water restrictions. Mice were randomly separated into four treatment groups ($n = 5$): (I)
177 vaccinated and infected with *B. abortus*, (II) vaccinated and non-infected with *B. abortus*, (III) non-
178 vaccinated and infected with *B. abortus*, and (IV) non-vaccinated and non-infected with *B. abortus*.
179 Mice were immunized intraperitoneally with a prime and two boosts (0, 7, 14 days) of vaccine
180 formulation containing 10 µg of peptide in phosphate-buffered saline (PBS), combined with
181 complete Freund's adjuvant (Sigma-Aldrich, St. Louis, MO, USA) at day 0 and incomplete
182 Freund's adjuvant at days 7 and 14. On day 21, mice were infected intraperitoneally with *B. abortus*
183 at a dose of 1×10^6 CFU/animal. Bacterial loads in the spleen, liver, and axillary lymph node from
184 individual animals were homogenized in PBS, serially diluted 10-fold, and plated on *Brucella* broth
185 agar (Difco, BD-Pharmingen, San Diego, CA). Plates were incubated at 37°C, and the CFU were
186 counted after 3 days as previously described (59). The experimental design is represented in Figure
187 2.
188
189

2.9 Histopathology and immunohistochemistry assays

190 The medial lobes of the mice liver were collected, fixed in 10% buffered formaldehyde solution,
191 dehydrated, diaphanized, and embedded in paraffin. Four-micrometer-thick tissue sections were
192 stained with hematoxylin and eosin (H&E). Digital images were captured and digitized with the
193 AxioVision LE software (Carl Zeiss, Oberkochen, Germany); all of the sample fields were
194 photographed for histopathological evaluation. The histopathological changes were analyzed by
195 Image Pro-PlusR 4.5 software (Media Cybernetics Inc., Silver Spring, MD, USA). The total number
196 and size of granulomas present in histological liver sections was determined using an Axiophot
197

198 microscope (Carl Zeiss, Oberkochen, Germany) with a 40x objective lens. Immunohistochemistry
199 was performed as previously described (60). Briefly, liver sections were hydrated and incubated
200 with 10% hydrogen peroxide in PBS for 30 min. After being washed with PBS, slides were
201 transferred to a humid chamber at room temperature, incubated with 25 mg/ml of skim milk for 45
202 min, and then incubated with a primary antibody for 30 min. For immunolabeling, diluted (1:5,000)
203 serum from a rabbit experimentally inoculated with *B. abortus* S19 strain was used as polyclonal
204 anti-*B. abortus* antibody. Then, tissue sections were washed with PBS, incubated with secondary
205 antibody for 20 min, washed again with PBS, and incubated for 20 min with streptavidin-peroxidase
206 from a commercial kit (LSAB + kit; Dako Corporation, Carpinteria, CA). The reaction was revealed
207 using 0.024% diaminobenzidine (DAB; Sigma), and sections were counterstained with Mayer's
208 hematoxylin.

209

210 **2.10 Measurement of NO into splenocyte culture supernatants**

211 Splens cells from C57BL/6 mice under treatment obtained after maceration were treated with ACK
212 buffer (0.15 M NH₄Cl, 1.0 mM KHCO₃, 0.1 mM Na₂EDTA, pH 7.2) to lyse red blood cells. After
213 that, the cells were washed with saline (NaCl 0.8%, wt/vol) and suspended in RPMI 1640 (Gibco,
214 Carlsbad, Calif) supplemented with 2 mM L-Glutamine, 25 mM HEPES, 10% (vol/vol) heat-
215 inactivated FBS (Gibco, Carlsbad, Calif), penicillin G sodium (100 U/mL), and streptomycin
216 sulfate (100 µg/mL). *Spleen* cells (1x10⁶) were cultured in 200µL *culture* medium and incubated at
217 37°C with 5% CO₂. The supernatant of splenocyte cultures was collected after 24h and nitric oxide
218 (NO) measurement was performed according to the Griess method (61).

219

220 **2.11 Analysis of surface markers CD86 and CD11b by fluorescence microscopic**

221 BMDMs were plated on imaging slides (µ-Slide 12-well, glass bottom, Ibidi GmbH, Munich,
222 Germany), followed by stimulation with splenocytes supernatant. The cells were then washed three
223 times with PBS and incubated with the anti-CD16/32 antibody (BD Biosciences, San Jose, CA) for
224 2 hours to block nonspecific bonds. The cells were then incubated with anti-CD86 and anti-CD11b,
225 followed by staining with FITC-conjugated and PE-conjugated (BD Biosciences), respectively,
226 overnight at 4°C. The slides were washed with PBS and the nuclei were stained with 150 ng/mL
227 40,6-diamino-2-phenylindole (DAPI; Thermo Scientific) for 1 hour. All images were captured
228 using a Nikon Eclipse 80i fluorescence microscope (Melville, New York, U.S.A). Image J software
229 was used to analyze the markings obtained for the nucleus (blue fluorescence), CD11b+ cells (green
230 fluorescence), and CD86+ cells (red fluorescence).

231

232 **2.12 Real-time RT-PCR for pro and anti-inflammatory cytokines expression**

233 Liver and spleen macerate as well as BMDMs stimulated with splenocytes supernatant from the
234 four experimental groups were homogenized with TRIzol reagent (Invitrogen) to isolate total RNA.
235 Reverse-transcription of 1 µg total RNA was performed using Illustra™ Ready-To-Go RT-PCR
236 Beads (GE Healthcare, Buckinghamshire, UK). Real-time RT-PCR was conducted in a final
237 volume of 10 µL containing the following: SYBR® Green PCR Master Mix (Applied Biosystems,
238 Foster City, CA, USA), with cDNA as the PCR template and primers to amplify specific fragments
239 corresponding to specific gene targets (Table 1). The PCR reaction was performed with ABI 7500
240 Real-Time PCR System (Applied Biosystems, Foster City, CA), using the following cycling
241 parameters: 60°C for 10 min, 95°C for 10 min, 40 cycles of 95°C for 15 sec and 60°C for 1 min,
242 and a dissociation stage of 95°C for 15 sec, 60°C for 1 min, 95°C for 15 sec, 60°C for 15 sec. All
243 data are presented as relative expression units after normalization to the β-actin gene. PCR
244 measurements were conducted in triplicate. The differences in the relative expression were
245 analyzed by analysis of variance (ANOVA) followed by Tukey's test (p < 0.05 denotes statistical
246 significance).

247

248 **2.13 Statistical analysis**

249 Graphs were created and data analysis was performed using GraphPad Prism 8 software (San
250 Diego, CA, USA), using one-way ANOVA or two-way ANOVA (Bonferroni *post hoc* test). Values
251 <0.05 were considered statistically significant.

252 **3 Results**

253 **3.1 High-throughput sequencing identifies the expression of small RNAs of *B. abortus* during** 254 **infection in macrophages**

255 In the small RNA libraries from infected macrophages, we observed that 7.26% of all mapped
256 sequences belonged to the *B. abortus* genome (Table 2). Knowing this, using the Bowtie software,
257 it was seen that the small RNAs were found to be distributed in both chromosomes of the bacteria.
258 By analyzing the depth of coverage of the sequences, we identified a total of 3954 regions of broad
259 mapping of small RNAs in the genome of *B. abortus*, 2694 in chromosome I and 1260 in
260 chromosome II. However, in local data of the bacterial genome, the presence of three peaks
261 (hotspots) was observed, with two in chromosome I and one in chromosome II, constituted by small
262 RNAs being mapped in the same position of the genome and large quantities (Supplementary Figure
263 1). With that, these three hotspots were selected and we next concentrated our analysis on selecting
264 the small RNAs that were mapped multiple times in these hotspots. In total, we identified one
265 hundred small RNAs with this characteristic which were then submitted to further analysis to detect
266 the target mRNA in NCBI.
267

268 **3.2 Apolipoprotein N-acyltransferase (BAB1_2158) is a putative target of *B. abortus* sRNAs** 269 **and its expression is diminished during earlier time of BMDM infection**

270 After the sequences were obtained from NCBI, the surface-associated proteins were selected by our
271 SCL prediction tools. In this phase, 10 proteins were included, making up the final list of surface-
272 associated proteins. The exposure of extracellular structures is an attraction for the immune system
273 in the recognition of antigens and, therefore, the topology predictive tools selected seven proteins
274 composed of extracellular domain amino acids. All selected extracellular portions were composed
275 of at least 35 amino acids. To gain more insight into the biological functions of these 7 proteins, an
276 analysis was performed to identify biological patterns associated with antigens. Of the 7 proteins
277 analyzed, all were inferred by homology in UniProtKB, and a few are those reviewed regarding
278 their biological function (Table 3). The information obtained by the Gene Ontology (GO) project
279 showed that only two of these proteins would be involved in the biosynthesis process of the bacterial
280 structure, yet one of them had a smaller number of exposed amino acids. The other, in addition to
281 exposing 277 amino acids in the extracellular portion, also participates in the bacterial
282 lipopolysaccharide biosynthesis process, becoming the target protein for further analyses. The other
283 proteins analyzed had a transmembrane transport function, in addition to mediating cellular vesicle
284 fusion processes.

285 Therefore, apolipoprotein N-acyltransferase (BAB1_2158) was selected for the further
286 analysis of differential expression during BMDM infection. Knowing that small RNAs were
287 expressed in large quantities by the bacterium during intracellular infection by *Brucella*, we
288 investigated the levels of mRNA expression of Int in samples of intracellular and extracellular
289 growth in a model of BMDM infection with *B. abortus*. The result of differential expression
290 analysis showed that during BMDM infection, the bacteria decreased the Int gene expression when
291 compared to an exponential extracellular growth model (Figure 3). Taking into account the results
292 of the *in silico* analysis, the negative expression of the Int coding gene may be related to a post-
293 transcriptional gene regulation process used by the bacteria to repress the expression of this protein
294 during intracellular infection, thus establishing its replicative niche in the host cell.

295 Since the results indicate a possible mechanism of gene expression control by the bacterium
296 in stressful intracellular environments, we were searching for exposed immunogenic epitopes in the
297 extracellular amino acid sequence of Int. The results obtained indicated approximately 60 epitopes
298 that showed a strong binding affinity for HLA molecules (HLA-DRB1 0101, HLA-DRB1 0301,
299 HLA-DRB1 0401, HLA-DRB1 0701, HLA-DRB1 1101, and HLA-DRB1 1501). In this stage, the

300 predictor tools selected epitopes composed of exactly 15 amino acids. When subjected to alignment
301 with the extracellular portion of the protein, the most promiscuous epitopes formed a conserved
302 region; from this region, the epitope "AIPYILESTPQALAH" was selected. To identify the position
303 of the selected putative immunogenic epitope in the Int, we performed molecular modeling,
304 showing the selected epitope highlighted in red with transmembrane helices indicated in the model
305 in green, as indicated in Figure 4. When submitted to *in silico* screening tests, the epitope was
306 shown to be antigenic in addition to showing 100% similarity with sequences present only in the
307 apolipoprotein N-acyltransferase of *B. abortus*. To increase the confidence of using the selected
308 epitope in future *in vitro* and *in vivo* evaluations, allergenicity was tested using the AlgPred
309 software, characterizing it as non-allergenic. Although when analyzing the physicochemical
310 properties, the epitope was shown to be unstable, since it did not reach the minimum cut-off value
311 for stability, it had a relatively good half-life in mammalian cells (Table 4); that is, despite its
312 instability in the medium, it is suggestive that when it binds to the MHCII cleft, it can present the
313 desired stability.

314

315 **3.3 Molecular docking between the selected putative immunogenic epitope from Int and** 316 **MHC-II shows high probabilities of interaction between them**

317 Although not all structures of MHC-II were found in the database (IEDB), we used the structure of
318 3 available proteins: HLA-DRB1:0101 (PDB: 5NI9), HLA-DRB1:0401 (PDB: 5V4M) and HLA-
319 DRB1:1501 (PDB: 5V4N). When performing the molecular docking of the
320 "AIPYILESTPQALAH" epitope, the results showed great interaction energy of all alleles based on
321 the AutoDock Vina software (Table 5). To represent the molecular docking between the epitope
322 and the MHC-II allele, we chose the interaction with the highest interaction power pointed out by
323 the software. The epitope in question interacted very well with the MHCII of the HLA-DRB1:0101
324 allele, having the best interaction values, with an energy of $-8.1 \text{ Kcal.mol}^{-1}$ and presenting a total
325 of 9 hydrogen bonds between the epitope and the MHC, with two bonds involving the amino acid
326 HIS259 of the MHC with the amino acids LEU303 at a distance of 3.09\AA and with the amino acid
327 ALA304 at a distance of 3.26\AA from the epitope. One link of amino acid ASN260 of MHC with
328 amino acid HIS305 was 3.21\AA from the epitope, one link of amino acid GLN242 of MHC with
329 amino acid PRO293 showed a distance of 3.09\AA from the epitope, two linkages involving amino
330 acid TYR238 of MHC with the amino acid ILE292 was 2.91\AA away and the link with the amino
331 acid ALA291 was at a distance of 3.00\AA from the epitope; also, a link of the amino acid LYS249
332 of the MHC with the amino acid GLU297 was at a distance of 3.06\AA and finally two bonds
333 involving amino acids TYR208 and HIS191 of the MHC, showed distances of 2.90\AA and 3.24\AA ,
334 respectively, with amino acid SER298 of the epitope (Figure 5A and B). In summary, the peptide
335 was shown to have a great ability to interact with MHC-II slits, especially with the HLA-
336 DRB1:0101 allele, showing excellent interaction with 9 of the 15 amino acids of the epitope.
337 Although minor binding strengths have been identified, this does not preclude the possibility of
338 interaction between peptide-alleles.

339

340 **3.4 Immunization of mice with epitope-based vaccine provides protects against *B. abortus*** 341 **infection**

342 After immunization, vaccinated and unvaccinated mice were challenged by intraperitoneal
343 infection with *B. abortus*; after seven days, organs were collected to determine bacterial loads. In
344 the spleen, liver and axillary lymph node of mice vaccinated with the peptide, fewer viable bacteria
345 were recovered when compared to the control group (Figure 6).

346 To evaluate the potential use of the epitope-based vaccine, the protection level induced in mice
347 against virulent challenge infection was assessed. The degree of vaccine efficacy in C57BL/6 mice
348 was determined by subtracting the mean CFU/organ recovered from mice after vaccination and
349 challenged from the mean CFU/organ recovered from non-vaccinated but challenged control mice.
350 At this time, it was seen that the presence of vaccinal peptide in the animal organism triggered a
351 higher degree of protection against infection, approximately 1.20/0.80/0.84-log in spleen, liver, and

352 axillary lymph node, respectively (Table 6). This result showed that the RV-selected epitope
353 provided significant protection to C57BL/6 mice against *B. abortus*.

354 355 **3.5 Unvaccinated mice showed greater liver damage when infected with *B. abortus***

356 *B. abortus* infection is associated with the formation of focal granulomatous lesions in the spleen,
357 liver, and lymphoid tissues of both humans and rodents, starting 1–2 weeks post-infection (62). To
358 determine the characteristics of liver pathology upon vaccination with peptide during *B. abortus*
359 infection, we performed the histopathological analysis of liver tissue from vaccinated and
360 unvaccinated C57BL/6 mice and, challenged with *B. abortus*. Infection with *B. abortus* resulted in
361 the formation of hepatic granulomas in both groups (Figure 7A and C). In morphometric analysis,
362 a greater number of granulomas was observed in the tissue of animals that were not vaccinated with
363 the peptide and challenged by bacteria compared to the vaccinated group (Figure 7G). The same
364 result was seen when analyzing the area, which was greater in granulomas from unvaccinated
365 animals (Figure 7H). Histopathological lesions during *Brucella* infection usually are associated
366 with the bacterial load. To determine the relationship between granuloma formation and the bacteria
367 present in the granuloma, we performed immunohistochemistry to immunolabel *B. abortus*. Figures
368 7E and 7F showed the presence of *B. abortus* in the granulomatous lesions presented in the liver of
369 vaccinated and infected mice and non-vaccinated and infected mice, respectively. The detection of
370 intralesional bacteria confirms that the inflammatory lesions described in this study are due to
371 systemic *B. abortus* infection. No observable lesions were found in tissues from uninfected mice
372 (Figure 7B and D).

373 374 **3.6 Vaccination induces positive expression of pro- and anti-inflammatory cytokines in** 375 **animals infected with *B. abortus***

376 The liver is the most commonly affected organ in patients with active brucellosis (63), which is
377 why liver macerates from the four experimental groups were collected for evaluation of the
378 expression of the anti-inflammatory genes *IL-10* and *TGF- β* . In the differential expression analysis,
379 an increase in these cytokines was observed in vaccinated and infected C57BL/6 mice when
380 compared to the other groups (Figure 8A-B). Therefore, the reduction in liver pathology can be
381 attributed to a decrease in the number of viable bacteria and an increase in anti-inflammatory
382 cytokines, resulting in a consequent reduction in liver damage. Concomitantly, the splenic tissue
383 was evaluated and the results obtained showed that all groups, except for the control, showed
384 upregulation of the proinflammatory cytokine-coding genes *INF- γ* , *TNF- α* , and *IL-6*, which are
385 characteristic of inflammation (Figure 8C-E), but the vaccinated and infected group stood out due
386 to the increased expression when compared to other groups. Likewise, it was seen that IL-10 (Figure
387 8E) is up-regulated in this group, suggesting that vaccination induces an attempt to control the
388 inflammatory process generated by systemic infection with *B. abortus*.

389 390 **3.7 BMDM's from C57BL/6 mice stimulated with supernatant splenocytes from the** 391 **vaccinated and infected group showed higher expression of CD86**

392 *In vivo* analysis results indicate the activation of an adaptive immune response. Knowing this, we
393 tried to understand the mechanisms by which a more efficient immune response activation process
394 against the bacteria occurs. BMDMs were evaluated for the expression of co-stimulatory molecules
395 after being stimulated with the splenocyte supernatant. The results obtained by fluorescence
396 microscopy show that BMDMs stimulated with the supernatant from the spleen of vaccinated and
397 infected animals were more activated, due to intense CD86 labeling (Figure 9A), and that they even
398 expressed a higher level of *iNOS* expression, the gene that stimulates NO production (Figure 9B).
399 To confirm this result, the measurement of NO was performed in the splenocyte supernatant used
400 to stimulate BMDMs. The results obtained showed the greater production of NO by supernatant
401 from the vaccinated and infected group, that is, a product that suggestively activated the BMDMs
402 more intensely (Figure 9C), which may reveal an increased phagocytic and microbicide capacity to
403 eliminate the bacteria.

404 **4 Discussion**

405 The development of subunit vaccines to protect against brucellosis is crucial to avoid the
406 disadvantages of the used live attenuated vaccines RB51 and S19 against *B. abortus* (64,65). New
407 vaccines will be designed according to immune responses during a natural infection in animal
408 models and the identification of intracellular and cell surface immunodominant components of
409 *Brucella* spp. (66-69). Using RV, it has been shown that the genome of *B. abortus* contains
410 approximately 80 genes encoding putative lipoproteins that have diverse functionalities, including
411 pathogenic processes (70,71). Knowing this, recent studies have shown that some bacterial cell
412 surface proteins can provide significant protection against *Brucella*, such as L7/L12 (72), Omp19,
413 Omp31 (73), BP26 (74), and Omp25 (75,76), which have been shown to be immunodominant
414 antigens that stimulate host immunity and trigger a protective response against infection in a mouse
415 model. Bacteria of the genus *Brucella* can live, replicate and persist within phagocytes using their
416 mechanisms to evade the immune system and establish their replicative niche within the host cell
417 (77). In this context, previous studies suggest that small RNAs may be directly related to the timely
418 gene expression of virulence factors in a variety of pathogenic bacteria such as *Listeria* and *Yersinia*
419 (78,79). Considering that the expression of small bacterial RNAs allows changes in the host cell
420 phenotype, and knowing that these small RNAs act on gene activation and repression, we evaluated
421 the capacity that a rationally predicted epitope of Int, a target protein of small RNAs expressed on
422 large scale by the bacteria during macrophage infection, would have to activate protective immune
423 responses during infection by *B. abortus* in a murine model.

424 Since *Brucella* species are equipped with a variety of well-organized immune evasion
425 strategies to establish chronic infections, including the use of small non-coding RNAs (80,81), we
426 performed a detailed analysis from data from the sequencing of small RNAs expressed during the
427 infection of BMDMs with *B. abortus*, showing that 7.26% of the small RNAs were mapped in the
428 bacterial genome. This was consistent with previous reports that identified expression levels of
429 similar small *B. abortus* RNAs during the infection of murine macrophages (82). Casewell et al.
430 reported that *B. abortus* small RNAs, abcR1 and abcR2 play essential roles in pathogenicity and
431 chronic infection, resulting in a significant decrease in intracellular survival in a mouse model and
432 in macrophages (83). Another group identified 129 small RNAs of *Brucella* that play significant
433 roles in diverse biological processes, ranging from physiology to virulence, as well as in host-
434 pathogen interaction (84). These reports shed light on the importance of small non-coding RNAs
435 in *Brucella* immunity, pathogenesis, and intracellular survival, modulating the host's immune
436 response. Here, we selected a non-allergenic but antigenic epitope with a good half-life in
437 mammals, yeast, or *E. coli*, from a putative target of a more highly expressed *Brucella* sRNA.
438 Rationally, we also take account of the structural and functional aspects of this target meeting the
439 epitope "AIPYILESTPQALAH" of apolipoprotein N-acyltransferase (Int). Although other proteins
440 were predicted as possible targets of *Brucella* sRNA in our analyses, this one, in particular, stood
441 out due to its biological function and strong epitope MHCII-interact capacity. Reportedly, Int
442 functionally constitutes *Brucella*'s outer membrane and plays a crucial role in bacterial LPS
443 biosynthesis (70,71,85). Interestingly, one of the main virulence factors of *Brucella* identified so
444 far is its non-canonical LPS (86,87) which exhibits favorable properties for the bacterium, including
445 low endotoxicity, high resistance to degradation in macrophages, and protection against immune
446 responses (88-90). The differential expression analysis of this protein-coding gene was performed
447 in intracellular growth samples in a BMDM model infected with *B. abortus*, showing a drastic
448 reduction in expression, corroborating our hypothesis that there is a post-transcriptional gene
449 regulation process used by the bacterium to repress the expression of Int during the early time of
450 infection, which can favor the bacteria permanence and replication in the host cell. Therefore, in
451 this study, in an unprecedented way, we evaluated the capacity of a specific Int epitope selected by
452 RV to induce an immune response in a murine model infected with *B. abortus*. The candidate
453 epitope-based vaccine was able to trigger protective immune responses, when the amount of viable
454 *B. abortus* in the liver, spleen, and axillary lymph nodes of vaccinated and unvaccinated mice when

455 challenged intraperitoneally by this pathogenic bacterium was evaluated. It was observed that
456 immunization considerably decreased the recovery of *B. abortus* in the tissues evaluated and
457 induced mean systemic protection of 0.94 logs when compared to unvaccinated animals. Other
458 evaluated *Brucella* antigens behaved similarly; for example, the recombinant Omp16 and Omp19
459 and the encapsulated recombinant liposome Omp25 induced protection comparable to S19 in
460 vaccinated mice after challenge (17,18,91). In addition, the Omp28 subunit vaccine increased
461 resistance against the *B. abortus* challenge by inducing a CD4+ Th1 response, that protects against
462 infection, but at a lower level than live attenuated vaccines (92). Corroborating our results, other
463 studies, when analyzing the level of splenic CFU, showed that the levels of protection in animals
464 immunized with RB51 averaged 0.91 logs [93,94], suggesting that the vaccine peptide in this study
465 induces levels of splenic protection similar to the live attenuated RB51 vaccine.

466 The characteristic pathological manifestation of *B. abortus* infection is granulomatous
467 inflammation associated with bacterial load (95). In this study, we detected a significant reduction
468 in the number and size of granulomas in the livers of animals vaccinated with the peptide compared
469 to unvaccinated animals, suggesting that the vaccination induced an effective inflammatory
470 immune response in this tissue. This condition is well defined, whereas, as infection in mice
471 infected with *Brucella* progresses, the granulomas progressively decrease in size and number after
472 2–3 weeks of infection (62). In parallel, we saw that the reduction of pathology in vaccinated mice
473 was accompanied by an increase in the expression of *IL-10* and *TGF- β* in the liver of infected
474 animals. Although the impact of IL-10 on *Brucella* persistence and the establishment of chronic
475 infection through macrophage modulation has been previously demonstrated using IL-10-deficient
476 mice (96,97), our findings provide evidence that the increased expression of *IL-10* is related to an
477 immunoregulatory mechanism dampening excessive Th1 responses (98). In this context, it is noted
478 that the absence of IL-10 results in severe pathological changes in different bacterial infections
479 (98,99). Here, we can speculate that the reduction in liver pathology in immunized and infected
480 animals may have been mediated by an increase in anti-inflammatory cytokines, and attributed to
481 a previous reduction in the number of viable bacteria.

482 Cell-mediated immunity is considered critical for the protective immune response against
483 facultative intracellular pathogens (100,101). Our results showed that the Int epitope induced the
484 enhanced production of TNF- α and IFN- γ in spleen cells, suggesting that there is some induction
485 of a Th1-type immune response by the vaccine peptide. In particular, IFN- γ is essential for immune
486 protection against *Brucella* infection that induces more polarization toward Th1 cells (102,103). In
487 addition, functional TNF- α has been shown to link the proinflammatory response and adaptative
488 immune response in *Brucella*-infected mice (104). High levels of IL-6 were produced by the
489 splenocytes of immunized and infected mice. It was recently shown that IL-6 is required for the
490 induction of IFN- γ and TNF- α by infected splenocytes, in addition to promoting the differentiation
491 of CD8+ T cells, indicating a protective role for IL-6 against *B. abortus* that parallels the type of
492 Th1 immunity response (105,106). Similarly, significant levels of IL-10 in immunized mice were
493 also detected in spleen cells. These results are consistent with the scenario seen in liver tissue,
494 suggesting that an inflammatory immune response has already occurred to the point of generating
495 an anti-inflammatory response with infection control characteristics. In this context, the balance
496 between the production of pro- and anti-inflammatory cytokines appears to be crucial for the host's
497 ability to eradicate the infection (97).

498 Since immunization triggered an effective adaptive immune response *in vivo*, the activation
499 of BMDMs when stimulated with supernatant from splenocytes from treated animals was evaluated
500 *in vitro*. It was observed that the supernatant from vaccinated and infected animals secreted cellular
501 components capable of stimulating macrophages more intensely in the BMDMs compared to the
502 other groups, even showing that these cells expressed higher levels of iNOS, and the consequent
503 increased production of NO, an important cell signaling molecule involved in infectious diseases
504 and the death of intracellular pathogens (107), justifying here the reduction in the number of viable
505 bacteria recovered from the vaccinated group. *In vivo*, IL-6, TNF- α , and CD80/CD86 are required
506 for activation of the interferon gamma-producing CD4+ Th1 and CD8+ cytotoxic T cells, a
507 protective response induced by the host against brucellosis (81,103,108,109). Although our study

508 did not assess the predominant subset of T cells in the immune response, it is believed that the
509 bactericidal and phagocytic function of macrophages to eliminate the bacteria was mainly enhanced
510 by the secretion of IFN- γ and TNF- α , since these cytokines showed high expression levels in tissues
511 from immunized animals, suggesting that the Int subunit vaccine predominantly induced an
512 effective Th1 profile response and triggered protection against *B. abortus* infection (Figure 10).

513 The information gathered shows that the bioinformatics is a strong approach for vaccine
514 candidate discovery as it offers a faster, cheaper, and safer method to identify potential vaccine
515 targets when compared with traditional laboratory identification methods, particularly when dealing
516 with risk group 3 microorganisms such as *Brucella*. Here, we provide an RV strategy that was able
517 to identify a *B. abortus* antigen that is found to be strongly associated with bacterial virulence.
518 Thus, immunization with the peptide vaccine had a significant effect on protection against murine
519 infection, inducing an immunoprotected response; therefore, it is plausible to assume that this
520 antigen can form a solid basis for designing an efficient and safe vaccine against animal brucellosis.

521 **6 Conflict of Interest**

522 The authors declare that the research was conducted in the absence of any commercial or financial
523 relationships that could be construed as a potential conflict of interest.

524 **7 Ethical Statement**

525 This study was conducted in strict accordance with the Brazilian laws 6638 and 9605 in Animal
526 Experimentation. The protocol was approved by the Committee on the Ethics of Animal
527 Experiments of the Federal University of Alfenas (CEUA 16/2020).

528 **8 Author Contributions**

529 KO and LA performed study design. RS, PC, and SO involved in contribution of study materials.
530 KO, GB, LA, and EN provided guidance for analytical tools and performed bioinformatic analysis.
531 KO and NS performed acquisition and collection of data *in vitro* and *in vivo*. KO and LA were
532 involved in manuscript preparation. All authors contributed to the article and approved the
533 submitted version.

534 **9 Funding**

535 This study was supported in part by the Coordenação de Aperfeiçoamento de Pessoal de Nível
536 Superior—Brasil (CAPES) (Finance Code 001) and Fundação de Amparo à Pesquisa do Estado de
537 Minas Gerais (Grant 864/14).

538 **10 References**

539 [1] Musallam II, Abo-Shehada MN, Hegazy, YM, Holt, HR, Guitian, FJ. (2016). Systematic review
540 of brucellosis in the Middle East: disease frequency in ruminants and humans and risk factors for
541 human infection. *Epidemiology and Infection*, 144, 671-685.
542 <https://doi.org/10.1017/S0950268815002575>.

543 [2] Khurana SK, Sehrawat A, Tiwari R, Prasad M, Gulati B, Shabbir MZ, et al. (2021). Bovine
544 brucellosis – a comprehensive review. *Veterinary Quarterly*, 41(1), 61-68.
545 <https://doi.org/10.1081/01652176.2020.1868616>.

546 [3] Pal M, Gizaw F, Fekadu G, Alemayehu G, Kandi V. (2017). Public health and economic
547 importance of bovine brucellosis: an overview. *American Journal of Epidemiology and Infectious
548 Disease*, 5(2), 27-34. <https://doi.org/10.12691/ajeid-5-2-2>.

- 549 [4] Gortázar C, Ferroglio E, Hofle U, Frolich K, Vicente J. (2007). Diseases shared between wildlife
550 and livestock a European perspective. *European Journal of Wildlife Research*, 53, 241-256.
551 <https://doi.org/10.1007/s10344-007-0098-y>.
- 552 [5] Franco MP, Mulder M, Gilman RH, Smits HL. (2007). Human brucellosis. *Lancet Infectious*
553 *Diseases*, 7(12), 775-786. [https://doi.org/10.1016/S1473-3099\(07\)70286-4](https://doi.org/10.1016/S1473-3099(07)70286-4).
- 554 [6] Ficht T. (2010). *Brucella* taxonomy and evolution. *Future Microbiology*, 5(6), 859-866.
555 <https://doi.org/10.2217/fmb.10.52>.
- 556 [7] Smits HL. (2013). Brucellosis in pastoral and confined livestock: prevention and vaccination.
557 *Revue Scientifique et Technique*, 32(1), 219, 228. <https://doi.org/10.20506/rst.32.1.2200>.
- 558 [8] Seleem MN, Boyle SM, Sriranganathan N. (2010). Brucellosis: a re-emerging zoonosis.
559 *Veterinary Microbiology*, 140(3-4), 392-398. <https://doi.org/10.1016/j.vetmic.2009.06.021>.
- 560 [9] Dadar M, Shahali Y, Fakhri Y, Godfroid J. (2020). The global epidemiology of *Brucella*
561 infections in terrestrial wildlife: A meta-analysis. *Transboundary and Emerging Diseases*,
562 2020(00), 1-15. <https://doi.org/10.1111/tbed.13735>.
- 563 [10] Franc KA, Krecek RC, Hasler BN, Arenas-Gamboa AM. (2018). Brucellosis remains a
564 neglected disease in the developing world: a call for interdisciplinary action. *BMC Public Health*,
565 18(125), 1-9. <https://doi.org/10.1186/s12889-017-5016-y>.
- 566 [11] Pappas G, Papadimitriou P, Akritidis N, Christou L, Tsianos EV. (2006). The new global map
567 of human brucellosis. *Lancet Infectious Diseases*, 6, 91-99. [https://doi.org/10.1016/s1437-3099\(06\)70382-6](https://doi.org/10.1016/s1437-3099(06)70382-6).
- 569 [12] Hisham Y, Ashhab Y. (2018). Identification of cross-protective potential antigens against
570 pathogenic *Brucella* spp. Through combining pan-genome analysis with reverse vaccinology.
571 *Journal of Immunology Research*, 2018(1474517), 1-15. <https://doi.org/10.1155/2018/1474517>.
- 572 [13] Khan MZ, Zahoor M. (2018). An overview of brucellosis in cattle and humans, and its
573 serological and molecular diagnosis in control strategies. *Tropical Medicine and Infectious Disease*,
574 3(2), 65. <https://doi.org/10.3390/tropicalmed3020065>.
- 575 [14] Adone R, Ciuchini F, Marianelli C, Tarantino M, Pistoia C, Marcon G, et al. (2005). Protective
576 properties of rifampin-resistant rough mutants of *Brucella melitensis*. *Infection and Immunity*,
577 73(7), 4198-4204. <https://doi.org/10.1128/IAI.73.7.4198-4204.2005>.
- 578 [15] Goodwin ZI., Pascual DW. (2016). Brucellosis vaccines for livestock. *Veterinary Immunology*
579 *and Immunopathology*, 181, 51-58. <https://doi.org/10.1016/j.vetimm.2016.03.011>.
- 580 [16] Saxena HM, Raj S. (2018). A novel immunotherapy of brucellosis in cows monitored non-
581 invasively through a specific biomarker. *PLoS Neglected Tropical Diseases*, 12(4),
582 e0006393. <https://doi.org/10.1371/journal.pntd.0006393>.
- 583 [17] Pasquevich KA., Estein SM, Samartino CG, Zwerdling A, Coria LM, Barrionuevo P, et al.
584 (2009). Immunization with recombinant *Brucella* species outer membrane protein Omp16 or
585 Omp19 in adjuvant induces specific CD4+ and CD8+ T cells as well as systemic and oral protection
586 against *Brucella abortus* infection. *Infection and Immunity*, 77(1), 436-445.
587 <https://doi.org/10.1128/IAI.01151-08>.

- 588 [18] Goel D, Rajendran V, Ghosh PC, Bhatnagar R. (2013). Cell mediated immune response after
589 challenge in Omp25 liposome immunized mice contributes to protection against virulent *Brucella*
590 *abortus* 544. *Vaccine*, 31(8), 1231-1237. <https://doi.org/10.1016/j.vaccine.2012.12.043>.
- 591 [19] Mallick AI, Singha H, Chaudhuri P, Nadeem A, Khan SA, Dar KA, et al. (2007). Liposomised
592 recombinant ribosomal L7/L12 protein protects BALB/c mice against *Brucella abortus* 544
593 infections. *Vaccine*, 25, 3692-3704. <https://doi.org/10.1016/j.vaccine.2007.01.066>.
- 594 [20] Grupta S, Mogan S, Somani VK, Aggarwal S, Bhatnagar R. (2020). Simultaneous
595 immunization with Omp25 and L7/L12 provides protection against brucellosis in mice. *Pathogens*,
596 9(152), 1-12. <https://doi.org/10.3390/pathogens9020152>.
- 597 [21] Al-Mariri A, Mahmoud NH, Hammoud R. (2012). Efficacy evaluation of live *Escherichia coli*
598 expression *Brucella* P39 protein combined with CpG oligodeoxynucleotides vaccine against
599 *Brucella melitensis* 16M, in BALB/c mice. *Biologicals*, 40(2012), 140-145.
600 <https://doi.org/10.1016/j.biologicals.2012.01.002>.
- 601 [22] Hop HT, Arayan LT, Huy TXN, Reyes AWB, Min W, Lee HJ, et al. (2018). Immunization of
602 BALB/c mice with a combination of four recombinant *Brucella abortus* proteins, AspC, Dps, InpB
603 and Ndk, confers a marked protection against a virulent strain of *Brucella abortus*. *Vaccine*, 36,
604 3027-3033. <https://doi.org/10.1016/j.vaccine.2018.04.019>.
- 605 [23] Ghasemi A, Jeddi-Tehrani M, Mautner J, Salari MH, Zarnani AH. (2015). Simultaneous
606 immunization of mice with Omp31 and TF provides protection against *Brucella melitensis*
607 infection. *Vaccine*, 33, 5532-5538. <https://doi.org/10.1016/j.vaccine.2015.09.013>.
- 608 [24] Wang X, Na C, Yang M, Li X, Ke Y, Lei S, et al. (2015). Immunization with individual
609 proteins of the Lrp/AsnC family induces protection against *Brucella melitensis* 16M challenges in
610 mice. *Frontiers in Immunology*, 6(1193), 1-8. <https://doi.org/10.3389/fmicb.2015.01193>.
- 611 [25] Carvalho TF, Haddad JPA, Paixão TA, Santos RL. (2016). Meta-analysis and advancement of
612 brucellosis vaccinology. *Plos One*, 11: e0166582. <https://doi.org/10.1371/journal.pone.0166582>.
- 613 [26] Bao Y, Tian M, Li P, Liu J, Ding C, Yu S. (2017). Characterization of *Brucella abortus* mutant
614 strain Δ 22915, a potential vaccine candidate. *Veterinary Research*, 48(1), 1-13.
615 <https://doi.org/10.1186/s13567-017-0422-9>.
- 616 [27] Rappuoli, R. (2000). Reverse vaccinology. *Current Opinion in Microbiology*, 3(5), 445-450.
617 [https://doi.org/10.1016/S1369-5274\(00\)00119-](https://doi.org/10.1016/S1369-5274(00)00119-).
- 618 [28] Pizza M, Scarlato V, Maignani V, Giuliani MM, Aricò B, Comanducci M, et al. (2000).
619 Identification of vaccine candidates against serogroup B *Meningococcus* by whole-genome
620 sequencing. *Science*, 287(5459), 1816-1820. <https://doi.org/10.1126/science.287.5459.1816>.
- 621 [29] Aslam M, Shehroz M, Shah M, Khan MA, Afridi SG, Khan A. (2020). Potential druggable
622 proteins and chimeric vaccine construct prioritization against *Brucella melitensis* from species core
623 genome data. *Genomics*, 112(2), 1734-1745. <https://doi.org/10.1016/j.ygeno.2019.10.009>.
- 624 [30] Seib KL, Zhao X, Rappuoli R. (2013). Developing vaccines in the era of genomics: a decade
625 of reverse vaccinology. *Clinical Microbiology and Infection*, 18(s5), 109-116.
626 <https://doi.org/10.1111/j.1469-0691.2012.03939.x>.

- 627 [31] Delany I, Rappuoli R, Seib KL. (2013). Vaccines, reverse vaccinology, and bacterial
628 pathogenesis. *Cold Spring Harbor Perspectives in Medicine*, 3(5), a012476.
629 <https://doi.org/10.1101/cshperspect.a012476>.
- 630 [32] Vishnu US, Sankarasubramanian J, Gunasekaran P, Rajendhran J. (2015). Novel vaccine
631 candidates against *Brucella melitensis* identified through reverse vaccinology approach. *OMICS:*
632 *A Journal of Integrative Biology*, 19(11), 722-729. <https://doi.org/10.1089/omi.2015.0105>.
- 633 [33] de Almeida LA, Carvalho NB, Oliveira FS, Lacerda TL, Vasconcelos AC, Nogueira L, et al.
634 (2011). MyD88 and STING signaling pathways are required for IRF3-mediated IFN- β induction in
635 response to *Brucella abortus* infection. *PloS One*, 6(8), e23135.
636 <https://doi.org/10.1371/journal.pone.0023135>.
- 637 [34] Martin M. (2011). Cutadapt removes adapter sequences from high-throughput sequencing
638 reads. *EMBnet Journal*, 17(1), 10-12. <http://dx.doi.org/10.14806/ej.17.1.200>.
- 639 [35] Bolger AM, Lohse M, Usadel B. (2014). Trimmomatic: a flexible trimmer of Illumina
640 sequence data. *Bioinformatics*, 30(15), 2114-2120. <https://doi.org/10.1093/bioinformatics/btu170>.
- 641 [36] Langmead B, Trapnell C, Pop M, Salzberg SL. (2009). Ultrafast and memory-efficient
642 alignment of short DNA sequences to the human genome. *Genome Biology*, 10(3), R25.1-R25.10.
643 <https://doi.org/10.1186/gb-2009-10-3-r25>.
- 644 [37] Quinlan AR, Hall IM. (2010). BEDTools: a flexible suite of utilities for comparing genomic
645 features. *Bioinformatics*, 26(6), 841-842. <https://doi.org/10.1093/bioinformatics/btp033>.
- 646 [38] Yu NY, Wagner JR, Laird MR, Melli G, Rey S, Lo R, et al. (2010). PSORTb 3.0: improved
647 protein subcellular localization prediction with refined localization subcategories and predictive
648 capabilities for all prokaryotes. *Bioinformatics*, 26(13), 1608-1615.
649 <https://doi.org/10.1093/bioinformatics/btq249>.
- 650 [39] Armenteros JJA, Tsirigos KD, Sonderby CK, Petersen TN, Winther O, Brunak S, et al. (2019).
651 SignalP 5.0 improves peptide predictions using deep neural networks. *Nature Biotechnology*, 37(4),
652 420-423. <https://doi.org/10.1038/s41587-019-0036-z>.
- 653 [40] Ikeda M, Arai M, Okuno T, Shimizu T. (2003). TMPDB: a database of experimentally-
654 characterized transmembrane topologies. *Nucleic Acids Research*, 31(1), 406-409.
655 <https://doi.org/10.1093/nar/gkg018>.
- 656 [41] Mitaku S, Hirokawa T, Tsuji T. (2002). Amphiphilicity index of polar amino acids as an aid
657 in the characterization of amino acid preference at membrane-water interfaces. *Bioinformatics*,
658 18(4), 608-616. <https://doi.org/10.1093/bioinformatics/18.4.608>.
- 659 [42] Krogh A, Larsson B, Heijne G, Sonnhammer ELL. (2001). Predicting transmembrane protein
660 topology with a hidden Markov model: application to complete genomes. *Journal of Molecular*
661 *Biology*, 305, 567-580. <https://doi.org/10.1006/jmbi.2000.4315>.
- 662 [43] Magrane M, Consortium U. (2011). UniProt Knowledgebase: a hub of integrated protein data.
663 *Database (Oxford)*, 2011, 1-13. <https://doi.org/10.1093/database/bar009>.
- 664 [44] Nielsen M, Lund O. (2009). NN-align. An artificial neural network-based alignment algorithm
665 for MHC class II peptide binding prediction. *BMC Bioinformatics*, 10(296), 1-10.
666 <https://doi.org/10.1186/1471-2105-10-296>.

- 667 [45] Rammensee HG, Bachmann J, Emmerich N, Bachor OA, Stevanovic S. (1999). SYFPEITHI:
668 database for MHC ligands and peptide motifs. *Immunogenetics*, 50, 213-219.
669 <https://doi.org/10.1007/s002510050595>.
- 670 [46] Reche PA, Glutting JP, Zhang H, Reinherz EL. (2004). Enhancement to the RANKPEP
671 resource for the prediction of peptide binding to MHC molecules using profiles. *Immunogenetics*,
672 56, 405-419. <https://doi.org/10.1007/s00251-004-0709-7>.
- 673 [47] Corpet, F. (1988). Multiple sequence alignment with hierarchical clustering. *Nucleic Acids*
674 *Research*, 16(22), 10881-10890. <https://doi.org/10.1093/nar/16.22.10881>.
- 675 [48] Doytchinova IA, Flower DR. (2007). VaxiJen: a server for prediction of protective antigens,
676 tumor antigens and subunit vaccines. *BMC Bioinformatics*, 8(4),1-7. <https://doi.org/10.1186/1471-2105/8/4>.
- 677
- 678 [49] Saha S, Raghava GPS. (2006). AlgPred: prediction of allergenic proteins and mapping of IgE
679 epitopes. *Nucleic Acids Research*, 34, W202-W209. <https://doi.org/10.1093/nar/gkl343>.
- 680 [50] Zai X, Yin Y, Guo F, Yang Q, Li R, Li Y, et al. (2021). Screening of potential vaccine
681 candidates against pathogenic *Brucella* spp. using composite reverse vaccinology. *Veterinary*
682 *Research*, 52(75), 1-15. <https://doi.org/10.1186/s13567-00939-5>.
- 683 [51] Gasteiger E, Gattiker A, Hoogland C, Ivanyi I, Appel RD, Bairoch A. (2003). ExPASy: The
684 proteomics server for in-depth protein knowledge and analysis. *Nucleic Acids Research*, 32(3),
685 3784-3788. <https://doi.org/10.1093/nar/gkg563>.
- 686 [52] Kelley LA, Mezulis S, Yates CM, Wass MN, Sternberg MJE. (2015). The Phyre2 web portal
687 for protein modeling, prediction and analysis. *Nature Protocols*, 10(6), 845-858.
688 <https://doi.org/10.1038/nprot.2015.053>.
- 689 [53] Sanner MF. (1999). Python: A programming language for software integration and
690 development. *Journal of Molecular Graphics and Modelling*, 17, 57-61.
- 691 [54] Ormo M, Cubitt AB, Kallio K, Gross LA, Tsien RY, Remington SJ. (1996). Crystal structure
692 of the *Aequorea victoria* green fluorescent protein. *Science*, 273, 1392-1395.
693 <https://doi.org/10.1126/science.273.5280.1392>.
- 694 [55] Trott O, Olson AJ. (2009). AutoDock Vina: Improving the speed and accuracy of docking with
695 a new scoring function, efficient optimization, and multithreading. *Journal of Computational*
696 *Chemistry*, 31, 455-461. <https://doi.org/10.1002/jcc.21334>.
- 697 [56] O'Boyle NM, Banck M, James CA, Morley C, Vandermeersch T, Hutchison GR. (2011). Open
698 Babel: An open chemical toolbox. *Journal of Cheminformatics*, 3(33), 1-14.
699 <https://doi.org/10.1186/1758-2946-3-3>.
- 700 [57] Laskowski RA, Swindells MB. (2011). LigPlot+: Multiple ligand-protein interaction diagrams
701 for drug discovery. *Journal of Chemical Information and Modeling*, 51, 2778-2786.
702 <https://dx.doi.org/10.1021/ci200227u>.
- 703 [58] Rigsby R, Parker AB. (2016). Using the PyMOL application to reinforce visual understanding
704 of protein structure. *Biochemistry and Molecular Biology Education*, 44(5), 433-437.
705 <https://doi.org/10.1002/bmb.20966>.

- 706 [59] Gomes MT, Campos PC, Pereira GDS, Bartholomeu DC, Splitter G, Oliveira SC. (2016).
707 TLR9 is required for MAPK/NF- κ B activation but does not cooperate with TLR2 or TLR6 to
708 induce host resistance to *Brucella abortus*. *Journal of Leukocyte Biology*, 99(5), 771-780.
709 <https://doi.org/10.1189/jlb.4A0815-346R>.
- 710 [60] Xavier MN, Paixão TA, Poester FP, Lage AP, Santos RL. (2009). Pathological,
711 immunohistochemical and bacteriological study of tissues and milk of cows and fetuses
712 experimentally infected with *Brucella abortus*. *Journal of Comparative Pathology*, 140(2-3), 149-
713 157. <https://doi.org/10.1016/j.jcpa.2008.10.004>.
- 714 [61] Brandão AP, Oliveira FS, Carvalho NB, Vieira LQ, Azevedo V, Macedo GC, et al. (2012).
715 Host susceptibility to *Brucella abortus* infection is more pronounced in IFN- γ knockout than IL-
716 12/ β 2-microglobulin double-deficient mice. *Clinical and Developmental Immunology*, 2012,
717 589494. <https://doi.org/10.1155/2012/589494>.
- 718 [62] Stranahan LW, Khalaf OH, Garcia-Gonzalez DG, Arenas-Gamboa AM. (2019).
719 Characterization of *Brucella canis* infection in mice. *PloS one*, 14(6), e0218809.
720 <https://doi.org/10.1371/journal.pone.02118809>.
- 721 [63] Giambartolomei GH, Delpino MV. (2019). Immunopathogenesis of Hepatic
722 Brucellosis. *Frontiers in Cellular and Infection Microbiology*, 9(423), 1-9.
723 <https://dx.doi.org/10.3389%2Ffcimb.2019.00423>.
- 724 [64] Avila-Calderón ED, Lopez-Merino A, Sriranganathan N, Boyle SM, Contreras-Rodríguez A.
725 (2013). A history of the development of *Brucella* vaccines. *BioMed Research International*,
726 2013(743509), 1-8. <https://doi.org/10.1155/2013/743509>.
- 727 [65] Olsen SC. (2013). Recent developments in livestock and wildlife brucellosis vaccination.
728 *Revue Scientifique et Technique (International Office of Epizootics)*, 32(1), 207-217.
729 <https://doi.org/10.20506/rst.32.1.2201>.
- 730 [66] Schurig GG, Sriranganathan N, Corbel MJ. (2002). Brucellosis vaccines: past, present and
731 future. *Veterinary Microbiology*, 90(1-4), 479-496. [https://doi.org/10.1016/S0378-1135\(02\)00255-9](https://doi.org/10.1016/S0378-1135(02)00255-9).
732 9.
- 733 [67] Yang X, Skyberg JA, Cao L, Clapp B, Thornburg T, Pascual DW. (2013). Progress in *Brucella*
734 vaccine development. *Frontiers in Biology*, 8(1), 60-77. <https://doi.org/10.1007/s11515-012-1196-0>.
735 0.
- 736 [68] Zhao Z, Yan F, Ji W, Luo D, Liu X, Xing L, et al. (2011). Identification of immunoreactive
737 proteins of *Brucella melitensis* by immunoproteomics. *Science China Life Sciences*, 54(9), 880-
738 887. <https://doi.org/10.1007/s11427-011-4218-2>.
- 739 [69] Yang Y, Wang L, Yin J, Wang X, Cheng S, Lang X, et al. (2011). Immunoproteomic analysis
740 of *Brucella melitensis* and identification of a new immunogenic candidate protein for the
741 development of brucellosis subunit vaccine. *Molecular Immunology*, 49(1-2), 175-184.
742 <https://doi.org/10.1016/j.molimm.2011.08.009>.
- 743 [70] Sternon JF, Godessart P, Gonçalves de Freitas R, Van der Henst M, Poncin K, Francis N, et
744 al. (2018). Transposon sequencing of *Brucella abortus* uncovers essential genes for growth in vitro
745 and inside macrophages. *Infection and Immunity*, 86(8), e00312-18.
746 <https://doi.org/10.1128/IAI.00312-18>.

- 747 [71] Goolab S, Roth RL, Van Heerden H, Crampton MC. (2015). Analyzing the molecular
748 mechanism of lipoprotein localization in *Brucella*. *Frontiers in Microbiology*, 6(1189), 1-20.
749 <https://doi.org/10.3389/fmicb.2015.01189>.
- 750 [72] Oliveira SC, Splitter GA. (1996). Immunization of mice with recombinant L7L12 ribosomal
751 protein confers protection against *Brucella abortus* infection. *Vaccine*, 14(10), 959-962.
752 [https://doi.org/10.1016/0264-410X\(96\)00018-7](https://doi.org/10.1016/0264-410X(96)00018-7).
- 753 [73] Cassataro J, Estein SM, Pasquevich KA, Velikovskiy CA., de la Barrera S, Bowden R, et al.
754 (2005). Vaccination with the recombinant *Brucella* outer membrane protein 31 or a derived 27-
755 amino-acid synthetic peptide elicits a CD4+ T helper 1 response that protects against *Brucella*
756 *melitensis* infection. *Infection and Immunity*, 73(12), 8079-8088.
757 <https://doi.org/10.1128/IAI.73.12.8079-8088.2005>.
- 758 [74] Yang X, Walters N, Robison A, Trunkle T, Pascual DW. (2007). Nasal immunization with
759 recombinant *Brucella melitensis* bp26 and trigger factor with cholera toxin reduces *B. melitensis*
760 colonization. *Vaccine*, 25(12), 2261-2268. <https://doi.org/10.1016/j.vaccine.2006.12.004>.
- 761 [75] Goel D, Bhatnagar R. (2012). Intradermal immunization with outer membrane protein 25
762 protects Balb/c mice from virulent *Brucella abortus* 544. *Molecular Immunology*, 51(2), 159-168.
763 <https://doi.org/10.1016/j.molimm.2012.02.126>.
- 764 [76] Paul S, Peddayelachagiri BV, Nagaraj S, Kingston JJ, Batra HV. (2018). Recombinant outer
765 membrane protein 25c from *Brucella abortus* induces Th1 and Th2 mediated protection against
766 *Brucella abortus* infection in mouse model. *Molecular Immunology*, 99, 9-18.
767 <https://doi.org/10.1016/j.molimm.2018.04.002>.
- 768 [77] Huy TXN, Nguyen TT, Reyes AWB, Vu SH, Min W, Lee HJ, et al. (2021). Immunization
769 With a Combination of Four Recombinant *Brucella abortus* Proteins Omp16, Omp19, Omp28, and
770 L7/L12 Induces T Helper 1 Immune Response Against Virulent *B. abortus* 544 Infection in
771 BALB/c Mice. *Frontiers in Veterinary Science*, 7, 1221.
772 <https://doi.org/10.3389/fvets.2020.577026>.
- 773 [78] Mraheil MA, Billion A, Mohamed W, Mukherjee K, Kuenne C, Pischmarov J, et al. (2011).
774 The intracellular sRNA transcriptome of *Listeria monocytogenes* during growth in macrophages.
775 *Nucleic Acids Research*, 39(10), 4235-4248. <https://doi.org/10.1093/nar/gkr033>.
- 776 [79] Ko KY, Kim JW, Her M, Kang SI, Jung SC, Cho DH, et al. (2012). Immunogenic proteins of
777 *Brucella abortus* to minimize cross reactions in brucellosis diagnosis. *Veterinary Microbiology*,
778 156(3-4), 374-380. <https://doi.org/10.1016/j.vetmic.2011.11.011>.
- 779 [80] Hanna N, Ouahrani-Bettache S, Drake KL, Adams LG, Köhler S, Occhialini A. (2013). Global
780 Rsh-dependent transcription profile of *Brucella suis* during stringent response unravels adaptation
781 to nutrient starvation and cross-talk with other stress responses. *BMC Genomics*, 14(1), 1-16.
782 <https://doi.org/10.1186/1471-2164-14-459>.
- 783 [81] Ahmed W, Zheng K, Liu ZF. (2016). Establishment of chronic infection: *Brucella's* stealth
784 strategy. *Frontiers in Cellular and Infection Microbiology*, 6(30), 1-12.
785 <https://doi.org/10.3389/fcimb.2016.00030>.
- 786 [82] Corsetti PP, de Almeida LA, Gonçalves ANA, Gomes MTR, Guimarães ES, Marques JT, et
787 al. (2018). miR-181a-5p regulates TNF- α and miR-21a-5p influences gualynate-binding protein 5
788 and IL-10 expression in macrophages affecting host control of *Brucella abortus* infection. *Frontiers*
789 *in Immunology*, 9, 1331, 1-14. <https://doi.org/10.3389/fimmu.2018.01331>.

- 790 [83] Caswell CC, Gaines JM, Ciborowski P, Smith D, Borchers CH, Roux CM, et al. (2012).
791 Identification of two small regulatory RNAs linked to virulence in *Brucella abortus* 2308.
792 *Molecular Microbiology*, 85(2), 345-360. <https://doi.org/10.1111/j.1365-2958.2012.08117.x>.
- 793 [84] Dong H, Peng X, Wang N, Wu Q. (2014). Identification of novel sRNAs in *Brucella abortus*
794 2308. *FEMS Microbiology Letters*, 354(2), 119-125. <https://doi.org/10.1111/1574-6968.12433>.
- 795 [85] Herrou J, Willet JW, Fiebig A, Czyż DM, Cheng JX, Ultee E, et al. (2019). *Brucella*
796 periplasmic protein EipB is a molecular determinant of cell envelope integrity and virulence.
797 *Journal of Bacteriology*, 201(12), e00134-19. <https://doi.org/10.1128/JB.00134-19>.
- 798 [86] Lapaque N, Moriyon I, Moreno E, Gorvel JP. (2005). *Brucella* lipopolysaccharide acts as a
799 virulence factor. *Current Opinion in Microbiology*, 8(1), 60-66.
800 <https://doi.org/10.1016/j.mib.2004.12.003>.
- 801 [87] Cardoso PG, Macedo GC, Azevedo V, Oliveira SC. (2006). *Brucella* spp. noncanonical LPS:
802 structure, biosynthesis, and interaction with host immune system. *Microbial Cell Factories*, 5(1),
803 1-11. <https://doi.org/10.1186/1475-2859-5-13>.
- 804 [88] Forestier C, Moreno E, Pizarro-Cerda J, Gorvel JP. (1999). Lysosomal accumulation and
805 recycling of lipopolysaccharide to the cell surface of murine macrophages, an in vitro and in vivo
806 study. *The Journal of Immunology*, 162(11), 6784-6791.
- 807 [89] Moreno E, Berman DT, Boettcher LA. (1981). Biological activities of *Brucella abortus*
808 lipopolysaccharides. *Infection and Immunity*, 31(1), 362-370. [https://doi.org/10.1128/iai.31.1.362-](https://doi.org/10.1128/iai.31.1.362-370.1981)
809 [370.1981](https://doi.org/10.1128/iai.31.1.362-370.1981).
- 810 [90] Forestier C, Deleuil F, Lapaque N, Moreno E, Gorvel JP. (2000). *Brucella abortus*
811 lipopolysaccharide in murine peritoneal macrophages acts as a down-regulator of T cell activation.
812 *The Journal of Immunology*, 165(9), 5202-5210. <https://doi.org/10.4049/jimmunol.165.9.5202>.
- 813 [91] Pasquevich KA, Ibañez AE, Coria LM, García Samartino C, Estein SM, Zwerdling A, et al.
814 (2011). An oral vaccine based on U-Omp19 induces protection against *B. abortus* mucosal
815 challenge by inducing an adaptive IL-17 immune response in mice. *PloS one*, 6(1), e16203.
816 <https://doi.org/10.1371/journal.pone.0016203>.
- 817 [92] Lim JJ, Kim DH, Lee JJ, Kim DG, Min W, Lee HJ, et al. (2012). Protective effects of
818 recombinant *Brucella abortus* Omp28 against infection with a virulent strain of *Brucella abortus*
819 544 in mice. *Journal of Veterinary Science*, 13(3), 287-292.
820 <https://dx.doi.org/10.4142/jvs.2012.13.3.287>.
- 821 [93] Lacerda TLS, Cardoso PG, De Almeida LA, da Cunha Camargo ILB, Afonso DAF, Trant CC,
822 et al. (2010). Inactivation of formyltransferase (wbkC) gene generates a *Brucella abortus* rough
823 strain that is attenuated in macrophages and in mice. *Vaccine*, 28(34), 5627-
824 5634. <https://doi.org/10.1016/j.vaccine.2010.06.023>.
- 825 [94] Truong QL, Cho Y, Kim K, Park BK, Hahn TW. (2015). Booster vaccination with safe,
826 modified, live-attenuated mutants of *Brucella abortus* strain RB51 vaccine confers protective
827 immunity against virulent strains of *B. abortus* and *Brucella canis* in BALB/c mice. *Microbiology*,
828 *161*(11), 2137-2148. <https://doi.org/10.1099/mic.0.000170>.
- 829 [95] Enright FM, Araya LN, Elzer PH, Rowe GE, Winter AJ. (1990). Comparative histopathology
830 in BALB/c mice infected with virulent and attenuated strains of *Brucella abortus*. *Veterinary*

- 831 *Immunology and Immunopathology*, 26(2), 171-182. [https://doi.org/10.1016/0165-2427\(90\)90065-](https://doi.org/10.1016/0165-2427(90)90065-)
832 Z.
- 833 [96] Corsetti PP, de Almeida LA, Carvalho NB, Azevedo V, Silva TM, Teixeira HC, et al. (2013).
834 Lack of endogenous IL-10 enhances production of proinflammatory cytokines and leads to *Brucella*
835 *abortus* clearance in mice. *PLoS One*, 8(9), e74729. <https://doi.org/10.1371/journal.pone.0074729>.
- 836 [97] Xavier MN, Winter MG, Spees AM, Nguyen K, Atluri VL, Silva TM, et al. (2013). CD4+ T
837 cell-derived IL-10 promotes *Brucella abortus* persistence via modulation of macrophage function.
838 *PLoS Pathogens*, 9(6), e1003454. <https://doi.org/10.1371/journal.ppat.1003454>.
- 839 [98] Saraiva M, O'garra A. (2010). The regulation of IL-10 production by immune cells. *Nature*
840 *Reviews Immunology*, 10(3), 170-181. <https://doi.org/10.1038/nri2711>.
- 841 [99] Belo VA, Pereira JA, Souza SFD, de Lima Tana F, Pereira BP, de Oliveira Lopes D, et al.
842 (2021). The role of IL-10 in immune responses against *Pseudomonas aeruginosa* during acute lung
843 infection. *Cell and Tissue Research*, 383(3), 1123-1133. <https://doi.org/10.1007/s00441-020->
844 03308-4.
- 845 [100] Martirosyan A, Moreno E, Gorvel JP. (2011). An evolutionary strategy for a stealthy
846 intracellular *Brucella* pathogen. *Immunological Reviews*, 240(1), 211-234.
847 <https://doi.org/10.1111/j.1600-065X.2010.00982.x>.
- 848 [101] Durward M, Radhakrishnan G, Harms J, Bareiss C, Magnani D, Splitter GA. (2012). Active
849 evasion of CTL mediated killing and low quality responding CD8+ T cells contribute to persistence
850 of brucellosis. *PLoS one*, 7(4), e34925. <https://doi.org/10.1371/journal.pone.0034925>.
- 851 [102] Murphy EA, Asselan JS, Parent MA, Zou B, Baldwin CL. (2001). Interferon- γ is crucial for
852 surviving a *Brucella abortus* infection in both resistant C57BL/6 and susceptible BALB/c mice.
853 *Immunology*, 2001(103), 511-518. 10.1046 / j.1365-2567.2001.01258.x.
- 854 [103] Vitry MA, De Trez C, Goriely S, Dumoutier L, Akira S, Ryffel B, et al. (2012). Crucial role
855 of gamma interferon-producing CD4+ Th1 cells but dispensable function of CD8+ T cell, B cell,
856 Th2, and Th17 responses in the control of *Brucella melitensis* infection in mice. *Infection and*
857 *Immunity*, 80(12), 4271-4280. <https://doi.org/10.1128/IAI.00761-12>.
- 858 [104] Zhan Y, Cheers C. (1998). Control of IL-12 and IFN- γ production in response to live or dead
859 bacteria by TNF and other factors. *The Journal of Immunology*, 161(3), 1447-1453.
- 860 [105] Hop HT, Huy TXN, Reyes AWB, Arayan LT, Vu SH, Min W, et al. (2019). Interleukin 6
861 promotes *Brucella abortus* clearance by controlling bactericidal activity of macrophages and CD8+
862 T cell differentiation. *Infection and Immunity*, 87(11), e00431-19.
863 <https://doi.org/10.1128/IAI.00431-19>.
- 864 [106] Guimarães ES, Martins JM, Gomes MTR, Cerqueira DM, Oliveira SC. (2020). Lack of
865 Interleukin-6 Affects IFN- γ and TNF- α Production and Early In Vivo Control of *Brucella abortus*
866 Infection. *Pathogens*, 9(12), 1040. <https://doi.org/10.3390/pathogens9121040>.
- 867 [107] Hu H, Tian M, Li P, Guan X, Lian Z, Yin Y, et al. (2020). *Brucella* Infection Regulates
868 Thioredoxin-Interacting Protein Expression to Facilitate Intracellular Survival by Reducing the
869 Production of Nitric Oxide and Reactive Oxygen Species. *The Journal of Immunology*, 204(3), 632-
870 643. <https://doi.org/10.4049/jimmunol.1801550>.

871 [108] Skendros P, Boura, PJRST. (2013). Immunity to brucellosis. *Revue Scientifique et Technique*,
872 32(1), 137-147. <https://doi.org/10.20506/rst.32.1.2190>.

873 [109] Dorneles EM, Sriranganathan N, Lage AP. (2015). Recent advances in *Brucella abortus*
874 vaccines. *Veterinary Research*, 46(1), 1-10. <https://doi.org/10.1186/s13567-015-0199-7>.

875 **9 Supplementary Material**


876 The Supplementary Material for this article can be found online:

877 **10 Data Availability Statement**

878 All datasets generated for this study are included in the article/[Supplementary Material](#).


879

880 **FIGURE AND TABLES LEGENDS**




881

882 **Figure 1.** Reverse vaccinology protocol workflow applied in this study to select a vaccinal epitope candidate.
 883 The one hundred protein sequences were obtained from the NCBI and ten proteins were localized in the
 884 plasma membrane by the Psortb tool. The TMPRED, SOSUI, and TMHMM tools identified 7 proteins with
 885 extracellular amino acid domain, but when analyzing the biological functions in the UniProtKB, only one
 886 was selected. Approximately 60 epitopes from the extracellular portion showed strong binding by HLAs
 887 according to the tools NetMHCII, SYFPEITHI, and RANKPEP. Using the Multalin tool the alignment was
 888 performed and the most promiscuous candidate among the epitopes was selected. n= number of proteins.




889

890 **Figure 2.** Experimental design of in vivo analysis. C57BL/6 mice were vaccinated three times with intervals
 891 of seven days between doses, the first dose being in the presence of complete Freund's adjuvant and the other
 892 doses using the same incomplete adjuvant. Seven days after the last immunization the mice were challenged
 893 with the virulent strain 2308 of *B. abortus* and on day twenty-eight the animals were euthanized to obtain a
 894 spleen, liver and axillary lymph node.




895

896 **Figure 3.** Decreased in the gene expression of apolipoprotein N-acyltransferase is shown in an intracellular-
 897 growing model. Differential expression analysis of apolipoprotein N-acyltransferase in model of
 898 extracellular and intracellular growth was measured by real-time RT-PCR. Statistically significant
 899 differences relative to the extracellular exponential growth are represented by an asterisk (* $P < 0.05$).




900

901 **Figure 4.** Position identification of selected epitope in apolipoprotein N-acyltransferase 3D. Molecular
902 modeling by homology for the apolipoprotein N-acyltransferase. Protein is represented in white; the selected
903 epitope is highlighted in red, and the transmembrane helices are indicated in the models in green.




904

905 **Figure 5.** Interaction between the selected epitope from apolipoprotein N-acyltransferase and MHC-II allele.
 906 (A) 2D diagram of the best interaction between the HLA-DRB1:0101 allele and the epitope, representing
 907 above the dotted line epitope amino acids and below line MHCII amino acids and (B) 3D diagram of the
 908 interactions.




909

910 **Figure 6.** Protective efficacy after three rounds of immunization. For protection evaluation, vaccinated mice
 911 were challenged with 1×10^6 CFU/bacteria. One-week post-challenge, organs were collected to determine
 912 bacterial loads to assess protection efficacy. A smaller number of viable bacteria was identified in vaccinated
 913 mice, indicating great protection efficacy. Data points were individual values of CFU determinations ($n = 5$)
 914 and analyzed using a student's t test. $*P < 0.05$ relative to the non-vaccinated and infected group.






915

916 **Figure 7.** Histopathology analysis, immunohistochemistry, and morphometric of hepatic tissue of *B. abortus*
 917 infected C57BL/6 mice. (A) Representative of hematoxylin- and-eosin-stained sections of hepatic tissue
 918 from mice vaccinated and infected, (B) vaccinated and non-infected, (B) unvaccinated and infected, and (D)
 919 non-vaccinated and non-infected. Immunohistochemistry sections of hepatic tissue from mice vaccinated
 920 and infected (E) and non-vaccinated and infected (F) mice containing the *B. abortus* inside the granuloma.
 921 The graphs analyze the granulomas of liver tissue sessions that were sequentially captured in terms of
 922 number (G) and area (H). Statistically significant differences relative to the non-vaccinated group are
 923 represented by an asterisk (* $P < 0.05$). The arrows indicate the *B. abortus* within the granuloma. Scale bars:
 924 50 μm.



925


926 **Figure 8.** Liver and spleen showed increased expression of pro and anti-inflammatory cytokine in mice
 927 vaccinated and infected with *B. abortus*. Differential expression analysis of (A) IL-10 and (B) TGF- β from
 928 the liver tissue and (C) IFN- γ , (D) TNF- α , (E) IL-6, and (F) IL-10 from the spleen of C57BL/6 mice from
 929 the four experimental groups evaluated in this study. Transcript levels were measured by real-time RT-PCR.
 930 Error bars represent the mean \pm SD of samples assayed in triplicate. * $P < 0.05$ relative to the non-vaccinated
 931 and non-infected group. # $P < 0.05$ relative to the non-vaccinated and infected group. & $P < 0.05$ relative to
 932 the vaccinated and non-infected group.

A**B****C**

933

934 **Figure 9.** *In-vitro* analysis showed an increased phagocytic and microbicide capacity in BMDM's. (A)
 935 Fluorescence microscopy shows activation of BMDM's by the CD86 molecule after being stimulated with
 936 supernatant from the spleen of C57BL/6 animals. (B) Differential expression analysis of iNOS in stimulated
 937 BMDM's was measured by real-time RT-PCR and (C) NO dosage by splenocyte supernatant using Griess
 938 method. Error bars represent the mean \pm SD of samples assayed in triplicate. * $P < 0.05$ relative to the non-
 939 vaccinated and non-infected group. # $P < 0.05$ relative to the non-vaccinated and infected group. & $P < 0.05$
 940 relative to the vaccinated and non-infected group. Scale bars: 20 μm .

941



942

943 **Figure 10.** Model proposed in this study suggesting optimal efficacy of the rationally predicted vaccine
 944 peptide. After being immunized and challenged, C57BL/6 mice were more resistant to infection by *B.*
 945 *abortus*, with less systemic recovery of viable bacteria and reduced tissue damage, mediated by an anti-
 946 inflammatory response. When evaluating the immune response profile, the predominance of characteristic
 947 components of the Th1 profile was observed, such as IFN- γ and TNF- α , suggesting that the vaccine peptide
 948 stimulates a profile mediated by CD4⁺ Th1 lymphocytes to secrete specific components such as IFN- γ , that
 949 mediates the production of NO and acts directly on infected macrophages, enhancing their microbicidal and
 950 phagocytic capacity and controlling the spread of systemic infection by this bacterium. Created with
 951 [BioRender.com](https://www.biorender.com/).

952

953

Table 1. Primers used in this study.

Primer	Forward Sequence	Reverse Sequence
β -actina	5'-AGGTGTGCACCTTTTATTGGTCTCAA-3'	5'-TGTATGAAGGTTTGGTCTCCCT-3'
TNF- α	5'-CATCTTCTCAAAATTCGAGTGACA-3'	5'-TGGGAGTAGACAAGGTACAACCC-3'
IFN- γ	5'-TCTGGAGGAAGTGGCAAAG-3'	5'-TTCAGACTTCAAAGAGTCTGAGG-3'
IL-10	5'-GGTTGCCAAGCCTTATCGGA-3'	5'-ACCTGCTCCACTGCCTTGCT-3'
IL-6	5'-CCAGGTAGCTATGGTACTCCAGAA-3'	5'-GATGGATGCTACCAAACCTGGA-3'
TGF- β	5'-TGACGTCACTGGAGTTGTACGG-3'	5'-GGTTCATGTCATGGATGGTGC-3'
iNOS	5'-CAGCTGGGCTGTACAAACCTT-3'	5'-CATTGGAAGTGAAGCGTTTCG-3'

954

955

956

957

958

959

Table 2. Data obtained after mapping of reads.

	<i>B. abortus</i> infected BMDMs	%	Control noninfected BMDMs	%
Processed Reads	35.713.113		46.516.010	
Reads mapped in the <i>B. abortus</i> (S2308) genome	2.593.062	7.26%	17.269	0,04% of the total
Reads mapped in the <i>Mus musculus</i> (GRCm38) genome	33.120.051	92.74%	46.498.741	99,96% of mapped reads

960

961 **Table 3.** List of proteins predicted to have transmembrane helices and respective biological functions.

Protein ID (NCBI)	Length (aa)	Single-line annotation (NCBI)	Biological function (UniProtKB)	Gene
WP_002968965.1	386	Lipase	Pathogenesis and negative regulation of endosome organization, and vesicle fusion	BAB2_0654
WP_002964284.1	270	Phosphatidate cytidyltransferase	CDP-diacylglycerol biosynthetic process	BAB1_1179
WP_002965220.1	532	Apolipoprotein N- acyltransferase	Lipoprotein biosynthetic process	BAB1_2158
WP_002971227.1	430	Xanthine/uracil/vitamin C permease family	Transmembrane transporter activity	BAB2_0578
WP_002971267.1	510	Amino acid permease	Transmembrane transporter activity	BAB2_0864
WP_002966986.1	412	MFS transporter superfamily	Transmembrane transporter activity (carbohydrate)	BAB1_1882
WP_002964796.1	400	OPGC	Transmembrane transporter activity (sugar)	BAB1_1718

962

963 **Table 4.** Similarity, antigenicity, allergenicity, and physical-chemical properties of selected epitope.

Sequence: AIPYILESTPQALAH									
Similarity	Allergenicity	Antigenicity	MW ^a	pI ^b	Stability	Half-life Reticulocytes	Half-life Yeast	Half-life <i>E. coli</i>	IgE epitope
100%	No	0.7800	1623.87	5.24	40.27	4.4h	>20h	>10h	No

964 ^aMW, molecular weight; ^b isoelectric point.

965

966 **Table 5.** Molecular docking results according to AutoDock Vina software.

MHC-II	Connection Power (Kcal.mol ⁻¹)	Hbonds	Non-ligand residues involved in hydrophobic contacts MHC	Non-ligand residues involved in hydrophobic contacts epitope
5NI9	-8,1	9	11	5
5V4M	-6,6	5	8	6
5V4N	-7,1	7	10	3

967

968

969

970 **Table 6.** Protective efficacy conferred by vaccinal peptide against *B. abortus* infection.

Group (n = 5)	Log ₁₀ CFU (Spleen) ^a	Log Protection	Log ₁₀ CFU (Liver)	Log Protection	Log ₁₀ CFU (Lymph node)	Log Protection
Vaccinated	5.34 ± 0.28*	1.20	4.96 ± 0.34*	0.80	5.30 ± 0.19*	0.84
Non-vaccinated	6.54 ± 0.63	-	5.74 ± 0.13	-	6.13 ± 0.43	-


971 Protection units of vaccinated group are compared with that of non-vaccinated with Student's t-test, **P* < 0.05 is statistically
 972 significant.

973 ^a CFU, colony-forming units.

974

975

976



977

978 **Supplementary Figure 1.** Arrangement of hotspots along the genome of *B. abortus* bacteria expressed
 979 during macrophage infection. (A) On chromosome I of *B. abortus* there is the formation of two hotspots
 980 characterizing regions of intense mapping of small RNAs and (B) on chromosome II there is the formation
 981 of only one hotspot.

2 CONCLUSÃO

A abordagem *in silico* empregada neste estudo permitiu identificar de forma racional um epítipo potencialmente antigênico e a partir dele contruir uma vacina de subunidade bem determinada. A imunização com a vacina de peptídeo apresentou efeito protetor significativo contra a infecção murina por *B. abortus*, induzindo uma resposta imunoprotetora e reduzindo os danos em tecidos acometidos pela infecção. Embora não tenha sido avaliado o subconjunto de células T predominantes na resposta imune, os resultados obtidos frente ao perfil imunológico de animais vacinados e infectados, sugere que a vacina de subunidade de apoliproteína N-aciltransferase induziu predominantemente uma resposta de perfil Th1 eficaz e desencadeou proteção contra a infecção por *B. abortus*. Portanto, é plausível supor que esse antígeno possa se tornar um forte candidato vacinal para o desenvolvimento de uma vacina melhorada, segura e eficiente contra a brucelose animal.

REFERÊNCIAS

- Abkar, M., Lotfi, A. S., Amani, J., Eskandari, K., Ramandi, M. F., Salimian, J., ... & Koushki, H. Survey of Omp19 immunogenicity against *Brucella abortus* and *Brucella melitensis*: influence of nanoparticulation versus traditional immunization. *Veterinary Research Communications*, 39(4), 217-228, 2015.
- Baldi, P. C., & Giambartolomei, G. H. Pathogenesis and pathobiology of zoonotic brucellosis in humans. *Revue Scientifique et Technique (International Office of Epizootics)*, 32(1), 117-125, 2013.
- Byndloss, M. X., & Tsolis, R. M. *Brucella* spp. virulence factors and immunity. *Annual Review of Animal Biosciences*, 4, 111-127, 2016.
- Carvalho, T. F., Haddad, J. P. A., Paixao, T. A., & Santos, R. L. Meta-analysis and advancement of brucellosis vaccinology. *PloS One*, 11(11), e0166582, 2016.
- Carvalho, T. F., Haddad, J. P. A., Paixão, T. A., & Santos, R. L. Meta-analysis of brucellosis vaccinology in natural hosts. *Pesquisa Veterinária Brasileira*, 40, 604-613, 2020.
- Casalinuovo, F., Ciambrone, L., Cacia, A., & Rippa, P. Contamination of bovine, sheep and goat meat with *Brucella* spp. *Italian Journal of Food Safety*, 5(3), 2016.
- de Oliveira, M. M., Pereira, C. R., de Oliveira, I. R. C., Godfroid, J., Lage, A. P., & Dorneles, E. M. S. Efficacy of *Brucella abortus* S19 and RB51 vaccine strains: a systematic review and meta-analysis. *Transboundary and Emerging Diseases*, 1-19, 2021.
- Dorneles, E. M., Teixeira-Carvalho, A., Araújo, M. S., Sriranganathan, N., & Lage, A. P. Immune response triggered by *Brucella abortus* following infection or vaccination. *Vaccine*, 33(31), 3659-3666, 2015.
- Elfaki, M. G., Alaidan, A. A., & Al-Hokail, A. A. Host response to *Brucella* infection: review and future perspective. *The Journal of Infection in Developing Countries*, 9(07), 697-701, 2015.
- El-Sayed, A., & Awad, W. Brucellosis: Evolution and expected comeback. *International Journal of Veterinary Science and Medicine*, 6, S31-S35, 2018.
- Escalona, E., Sáez, D., & Oñate, A. Immunogenicity of a multi-epitope DNA vaccine encoding epitopes from Cu-Zn superoxide dismutase and open reading Frames of *Brucella abortus* in mice. *Frontiers in Immunology*, 8, 125, 2017.

Franc, K. A., Krecek, R. C., Häsler, B. N., & Arenas-Gamboa, A. M. Brucellosis remains a neglected disease in the developing world: a call for interdisciplinary action. *BMC Public Health*, 18(1), 1-9, 2018.

Ghasemi, A., Jeddi-Tehrani, M., Mautner, J., Salari, M. H., & Zarnani, A. H. Immunization of mice with a novel recombinant molecular chaperon confers protection against *Brucella melitensis* infection. *Vaccine*, 32(49), 6659-6666, 2014.

Głowacka, P., Żakowska, D., Naylor, K., Niemcewicz, M., & Bielawska-Drozd, A. *Brucella*-virulence factors, pathogenesis and treatment. *Polish Journal of Microbiology*, 67(2), 151, 2018.

Gomez, G., Adams, L. G., Ficht, A. R., & Ficht, T. A. Host-*Brucella* interactions and the *Brucella* genome as tools for subunit antigen discovery and immunization against brucellosis. *Frontiers in Cellular and Infection Microbiology*, 3, 17, 2013.

Gupta, S., Mohan, S., Somani, V. K., Aggarwal, S., & Bhatnagar, R. Simultaneous immunization with Omp25 and L7/L12 provides protection against Brucellosis in mice. *Pathogens*, 9(2), 152, 2020.

Gutiérrez-Jiménez, C., Hysenaj, L., Alfaro-Alarcón, A., Mora-Cartín, R., Arce-Gorvel, V., Moreno, E., ... & Barquero-Calvo, E. Persistence of *Brucella abortus* in the bone marrow of infected mice. *Journal of Immunology Research*, 2018.

Hasanjani, M. R., & Ebrahimpour, S. Human brucellosis: An overview. *Caspian J Intern Med*, (2015): 46-47.

Hull, N. C., & Schumaker, B. A. Comparisons of brucellosis between human and veterinary medicine. *Infection Ecology & Epidemiology*, 8(1), 1500846, 2018.

Jain, S., Afley, P., Dohre, S. K., Saxena, N., & Kumar, S. Evaluation of immunogenicity and protective efficacy of a plasmid DNA vaccine encoding ribosomal protein L9 of *Brucella abortus* in BALB/c mice. *Vaccine*, 32(35), 4537-4542, 2014.

Jamil, T., Melzer, F., Saqib, M., Shahzad, A., Khan Kasi, K., Hammad Hussain, M., ... & Neubauer, H. Serological and molecular detection of bovine brucellosis at institutional livestock farms in Punjab, Pakistan. *International Journal of Environmental Research and Public Health*, 17(4), 1412, 2020.

Jezi, F. M., Razavi, S., Mirnejad, R., & Zamani, K. Immunogenic and protective antigens of *Brucella* as vaccine candidates. *Comparative Immunology, Microbiology and Infectious Diseases*, 65, 29-36, 2019.

Lalsiamthara, J., & Lee, J. H. Development and trial of vaccines against *Brucella*. *Journal of Veterinary Science*, 18(S1), 281-290, 2017.

Lindahl-Rajala, E., Hoffman, T., Fretin, D., Godfroid, J., Sattarov, N., Boqvist, S., ... & Magnusson, U. Detection and characterization of *Brucella* spp. in bovine milk in small-scale urban and peri-urban farming in Tajikistan. *PLoS Neglected Tropical Diseases*, 11(3), e0005367, 2017.

Nol, P., Olsen, S. C., Rhyan, J. C., Sriranganathan, N., McCollum, M. P., Hennager, S. G., ... & Salman, M. D. Vaccination of elk (*Cervus canadensis*) with *Brucella abortus* strain RB51 overexpressing superoxide dismutase and glycosyltransferase genes does not induce adequate protection against experimental *Brucella abortus* challenge. *Frontiers in Cellular and Infection Microbiology*, 6, 10, 2016.

O'callaghan, D. Human brucellosis: recent advances and future challenges. *Infectious Diseases of Poverty*, 9(1), 1-2, 2020.

Olsen, S. C., McGill, J. L., Sacco, R. E., & Hennager, S. G. Immune responses of bison and efficacy after booster vaccination with *Brucella abortus* strain RB51. *Clinical and Vaccine Immunology*, 22(4), 440-447, 2015.

Pizza, M., Scarlato, V., Maignani, V., Giuliani, M. M., Arico, B., Comanducci, M., ... & Rappuoli, R. Identification of vaccine candidates against serogroup B *meningococcus* by whole-genome sequencing. *Science*, 287(5459), 1816-1820, 2000.

Poester, F. P., Samartino, L. E., & Santos, R. L. Pathogenesis and pathobiology of brucellosis in livestock. *Rev Sci Tech*, 32(1), 105-115, 2013.

Tana, F. L., Guimarães, E. S., Cerqueira, D. M., Campos, P. C., Gomes, M. T. R., Marinho, F. V., & Oliveira, S. C. Galectin-3 regulates proinflammatory cytokine function and favors *Brucella abortus* chronic replication in macrophages and mice. *Cellular Microbiology*, e13375, 2021.

Tian, M., Song, M., Yin, Y., Lian, Z., Li, Z., Hu, H., ... & Yu, S. Characterization of the main immunogenic proteins in *Brucella* infection for their application in diagnosis of brucellosis. *Comparative Immunology, Microbiology and Infectious Diseases*, 70, 101462, 2020.

Vishnu, U. S., Sankarasubramanian, J., Gunasekaran, P., & Rajendhran, J. Novel vaccine candidates against *Brucella melitensis* identified through reverse vaccinology approach. *Omic: A Journal of Integrative Biology*, 19(11), 722-729, 2015.